Write your name here	l ou	
Surname	Otne	er names
Edexcel International GCSE	Centre Number	Candidate Number
Chemistry Unit: 4CH0	y	
Science (Double Av Paper: 1C	vard) 4SC0	
Friday 13 January 2012 – Time: 2 hours	Morning	Paper Reference 4CH0/1C 4SC0/1C
You must have: Ruler Calculator.		Total Marks

## **Instructions**

- Use **black** ink or ball-point pen.
- **Fill in the boxes** at the top of this page with your name, centre number and candidate number.
- Answer **all** questions.
- Answer the questions in the spaces provided
  - there may be more space than you need.
- Show all the steps in any calculations and state the units.

## Information

- The total mark for this paper is 120.
- The marks for **each** question are shown in brackets
  - use this as a guide as to how much time to spend on each question.

## **Advice**

- Read each question carefully before you start to answer it.
- Keep an eye on the time.
- Write your answers neatly and in good English.
- Try to answer every question.
- Check your answers if you have time at the end.

Turn over ▶



L	ı	ı	
		֚֚֡֡֡	
1	٥	ζ	
(		)	
1		5	
(	_	2	
Č	I	]	
(	1	-	
Ļ	I	1	
ŀ		=	

Helium 2

0

7

9

2

က

Group

α

Period

N

က

1 H Hydrogen 1

1         8         11         12         14         15         14         15         18 <th>0</th> <th>•</th> <th>- To 6</th> <th></th> <th></th> <th>_</th> <th>no. 8</th> <th>4</th> <th></th> <th>oton</th> <th>9</th> <th></th> <th><b>.</b></th> <th>nor 4.</th> <th>83</th> <th></th> <th>rop 4</th> <th>,</th> <th></th> <th></th> <th></th> <th></th>	0	•	- To 6			_	no. 8	4		oton	9		<b>.</b>	nor 4.	83		rop 4	,				
Bachilum   Bachilum	26	Z	Š.	=	4	⋖	A B T	œ		Kry,	e	5	× 	× ×	22	Œ	Æ ª	°				
Pach	19	ш	Fluorine	6	35.5	ರ	Chlorine 17	98	ă	Bromine	32	127	_	lodine 53	210	¥	Astatine	3				
Ba	9	0	Oxygen	8	35	တ	Sulfur 16	79	Se	Selenium	86	128	e H	Tellurium 52	210	Ъ	Polonium 84	5				
BBOON         11         B Boon         11         B Boon         11         B Boon         12         14         B Boon         15         27         All Aluminium         Aluminium         All Aluminium         Aluminium<	41	z	Nitrogen	,	3	۵.	Phosphorus 15	75	As	Arsenic	33	122	Sp	Antimony 51	209	ö	Bismuth	3				
Be Beryllium         45         48         51         52         55         56         59         59         63.5         65           Calcium         Scandium         Titanium         Vanadium         Chromium         Manganesium         Fe         CO         Ni         CO         Sn           40         45         48         51         52         55         56         59         63.5         65         59           2actium         Scandium         Titanium         Vanadium         Chromium         Manganese         Iron         Cobalt         Nickel         Copper         Zin           2actium         2acadium         Variuum         Arium	12	ပ	Carbon	9	88	ίδ	Silicon 14	73	Ĝe	Germanium	35	119	S	Έ ያ	207	P	Lead	90				
Be Beryllium Aggnesium Scandium String         51         52         55         56         59         59         635           Ca Calcium Scandium String         Vanadium String         Chromium Varnadium Chromium Manganese Scott         Cobatt Scott         Nickel Copper Scott         Copatt Scott         Nickel Copper Scott           So Calcium Scandium Tilanium Varnadium String         Varnadium Chromium Manganese Scott         Fe CO Ni Copatt Scott         Nickel Copper Scott         Copper Scott           So Sc Ti Varium Sirconium Stronium String         String Scott         Nickel Scott         Copper Scott         Nickel Scott         Copper Scott           Stronium String         Vitrium Sirconium String         Nickel Scott         Nickel Scott         Copatt Scott         Nickel Scott         Copper Scott           Stronium String         Nickel Scott         Nickel Scott         Nickel Scott         Nickel Scott         Nickel Scott         Nickel Scott           Stronium String         Nickel Scott         Nickel Scott         Nickel Scott         Nickel Scott         Nickel Scott         Nickel Scott           Stronium String         Nickel Scott         Nickel Scott         Nickel Scott         Nickel Scott         Nickel Scott         Nickel Scott           Stronium String         Nickel Scott         Nickel Scott         Nickel Scott <td>F</td> <td>æ</td> <td>Boron</td> <td>2</td> <td>27</td> <td>₹</td> <td>Aluminium 13</td> <td>8</td> <td>Ga</td> <td>Gallium</td> <td>31</td> <td>115</td> <td>드</td> <td>Indium 49</td> <td>204</td> <td>F</td> <td>Thallium</td> <td>ō</td> <td></td> <td></td> <td></td> <td></td>	F	æ	Boron	2	27	₹	Aluminium 13	8	Ga	Gallium	31	115	드	Indium 49	204	F	Thallium	ō				
Beryllium         4         5         5         55         56         59         59         59           Adordium Angenesium 12         45         48         51         52         55         56         59         59         59         59         59         59         59         59         59         59         59         59         59         59         59         59         59         50         59         50         59         50         59         59         50         59         50         59         50         59         50         59         50         59         50         59         50         59         50         59         50         59         50         59         50         59         50								65	Zu	Zinc	30	112	පි	Cadmium 48	201	모	Mercury	00				
Beryllium         45         48         51         52         55         56         59           Ca         Sc         Ti         V         Cr         Mn         Fe         Co           Calcium         Scandium         Titanium         V         Cr         Mn         Fe         Co           Co         Sc         Ti         V         Cr         Mn         Fe         Co           So         31         93         96         99         101         103         103           Strontium         Y         Zr         Nb         Mo         TC         Ru         Rh         44         45           Strontium         Y         Zr         Nb         Mo         TC         Ru         44         45         26         27           Ba         By         179         181         Ta         W         Re         44         45         44         45         44         45         45         45         45         45         45         45         45         45         44         45         44         45         77         77         77         77         77         77         77								63.5	J	Copper	29	108	Ag	Silver 47	197	Au	Gold	6/				
Be         Be           Be         4           24         As           Magnesium         45           12         Ca           Ca         Sc           City         Cranium           Sondium         Titanium           V         Cr           Sr         Y           Srontium         Virium           Srontium         Virium           Srontium         Virium           Srontium         Virium           Srontium         Virium           Srontium         Virium           Sig         91           40         41           41         72           As         40           41         42           42         43           44         42           45         43           46         41           47         42           40         41           41         42           43         44           44         42           44         44           45         73      73         74      74 <td< td=""><td></td><td></td><td></td><td></td><td></td><td></td><td></td><td>59</td><td>Z</td><td>Nickel</td><td>28</td><td>106</td><td>Pd</td><td>Palladium 46</td><td>195</td><td>ā</td><td>Platinum</td><td>9/</td><td></td><td></td><td></td><td></td></td<>								59	Z	Nickel	28	106	Pd	Palladium 46	195	ā	Platinum	9/				
Be   Beryllium   As   Barum   Chromium   C								59	රි	Cobalt	27	103	뜐	Rhodium 45	192	_	Iridium	*				
Be   Beryllium   4   As   As   As   As   As   As   As								26	E.	ro	56	101	2	Ruthenium 44	95	ő	Osmium	0/				
Be   Beryllium   4   As   As   As   As   As   As   As								55	Ž	Manganese	52	8	ည	Technetium 43	186	Be	Rhenium	6				
9   9   9   9   9   9   9   9   9   9																>	Tungsten	4/				
Be Beryllium 4  Mg Magnesium 12  40  Ca Calcium Scandium 20  21  88  88  88  87  Strontium 38  137  139  Ba La La Barium Lanthanum 56  26  27  28  88  89  88  89  89  87  21  88  89  89  89  89  89  89  89  89  8								51	>	Vanadium	23	83	S	Niobium 41	181	E	Tantalum	23				
Be Pervillam Magnesium 12 Calcium 20 Sr Strontium 38 Ba Barium 137 Se								48	F	Titanium	22	91	Z	Zirconium 40	179	Ĭ	Hafnium	7/				
Li   Be     Lithium   3								45	S	Scandium	24	88	>	Yttrium 39	139	2	Lanthanum	5	227	Ac	Actinium	5
Lithium 33 Na Sodium 11 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1	o	Be	Beryllium	4	24	Ž	Magnesium 12	40	ő	Calcium	ଷ	88	ഗ്	Strontium 38	137	Ва	Barium	8	526	Ва	Radium	000
	7	=	Lithium	3		Z	Sodium 11	96	¥	Potassium	19	98	8	Rubidium 37	133	S	Caesium	8	223	Ŧ	Francium	-

Key

Relative atomic mass Symbol Name Atomic number

P 4 0 1 2 6 A 0 2 2 8

4

2

9


/

**BLANK PAGE** 



## Answer ALL questions.

1 Salt is soluble in water, but sand is insoluble in water. This difference allows a mixture of salt and sand to be separated using this apparatus.



(a) Use words from the box to complete the sentences. Each word may be used once, more than once or not at all.

**(6)** 

beaker	Bunsen burner	burette	conical flask
funnel	glass rod	thermometer	water

In **Step 1**, the mixture of salt and sand is placed in a

containing and stirred with a

In Step 2, the mixture from Step 1 is poured through a

into a .....

In Step 3, the liquid is transferred to a basin to allow the.....

to be removed.



(b) (i)	What should be placed in A before the mixture from Step 1 is poured through it	t? (1)
(ii)	What is the solid removed in <b>Step 2</b> ?	(1)
(c) Plac	ce crosses ( ) in <b>two</b> boxes to show the names of two processes used in this separate	aration.
X	chromatography	
X	condensation	
X	distillation	
X	evaporation	
X	filtration	
$\times$	sublimation	
	(Total for Question 1 = 10 man	rks)

2	2 Iron is a useful metal. One problem with using iron is that it can rust.							
	(a) (i)	Name the iron compound present in rust.	(1)					
1	(ii)	Name the <b>two</b> substances that iron reacts with when it rusts.	(2)					
2								
	(b) Wh	at type of reaction occurs in the rusting of iron?						
	Plac	ce a cross (⋈) in <b>one</b> box.	(1)					
	$\boxtimes$	combustion						
	$\times$	decomposition						
	$\boxtimes$	displacement						
	$\boxtimes$	oxidation						
		vanising can prevent iron from rusting. In this process, the iron is coated with						
	ano	ther metal.						
		Identify the other metal.	(1)					
	(i)		(1)					
	(i)	Identify the other metal.	(1)					
	(i)	Identify the other metal.  Identify one object suitable for galvanising.						
	(i) (ii)	Identify the other metal.  Identify one object suitable for galvanising.  Place a cross (⋈) in one box.						
	(i) (ii)	Identify the other metal.  Identify one object suitable for galvanising.  Place a cross (⋈) in one box.  bicycle chain						
	(i) (ii)	Identify the other metal.  Identify one object suitable for galvanising.  Place a cross (⋈) in one box.  bicycle chain bucket						
	(i) (ii)	Identify the other metal.  Identify one object suitable for galvanising.  Place a cross (⋈) in one box.  bicycle chain bucket car engine						
	(i) (ii)	Identify the other metal.  Identify one object suitable for galvanising.  Place a cross (⋈) in one box.  bicycle chain bucket car engine						
	(i) (ii)	Identify the other metal.  Identify one object suitable for galvanising.  Place a cross (⋈) in one box.  bicycle chain bucket car engine						



	(d) State <b>two</b> other methods used to prevent iron from rusting. (2)
1	
2	
_	(Total for Question 2 = 8 marks)




3	Ammonium chloride contains oppositely charged ions.	
	(a) State the formula of each ion.	(2)
Po	sitive ion	(2)
Ne	gative ion	
	(b) (i) Describe a chemical test to show that a substance contains ammonium ions.	(3)
	(ii) Describe a chemical test to show that a substance contains chloride ions.	(3)
	(c) Ammonium chloride decomposes when heated:	
	$NH_4Cl(s) \rightleftharpoons NH_3(g) + HCl(g)$	
	What does the	(1)

(d) The reaction between ammonia and hydrogen chloride can be used to illustrate diffusion
with the following apparatus.
cotton wool soaked in concentrated in concentrated hydrochloric acid white solid ammonia solution
After a few minutes, a white solid appears inside the tube.
(i) Identify the white solid. (1)
(ii) What does the diagram show about the speed of the ammonia molecules compared to the speed of the hydrogen chloride molecules?  (1)
(e) State the main hazard when using concentrated hydrochloric acid in the experiment in (d).  Suggest <b>one</b> precaution you could use to minimise this hazard.
(2)
Hazard
Precaution
(Total for Question 3 = 13 marks)



4 A student set up the following apparatus.



- (a) The reaction between magnesium and hydrochloric acid forms hydrogen gas.
  - (i) State **one** observation the student would make during this reaction.

(1)

(ii) Identify the other product formed during this reaction.

(1)

- (b) The hydrogen gas burns in air to form steam. The steam changes to water on the surface of the round flask.
  - (i) Write a chemical equation for the burning of hydrogen in air.

(2)

(ii) What name is used for the process in which steam changes into water?


(1)

(i) State the final colour of	of the copper(II) sulfate in this reaction.	(1)
(ii) The colour change o	f the anhydrous copper(II) sulfate shows th	at the liquid contains was
Describe a test to sho	ow that the water is pure.	(2)
	(Total for Que	estion 4 = 8 marks)
	(Total for Que	estion 4 = 8 marks)
	(Total for Qu	estion 4 = 8 marks)
	(Total for Qu	estion 4 = 8 marks)
	(Total for Qu	estion 4 = 8 marks)
	(Total for Que	estion 4 = 8 marks)
	(Total for Que	estion 4 = 8 marks)
	(Total for Que	estion 4 = 8 marks)
	(Total for Que	estion 4 = 8 marks)



5 These are the displayed formulae of six organic compounds.

H   H—C—H   H	H H	H H H
P	Q	R



- (a) Use the letters above to select
  - (i) the compound that is **not** a hydrocarbon.

(1)

(ii) one compound with the empirical formula CH<sub>2</sub>

(1)

(iii) one compound that can form a polymer.

(1)

(b) Describe a test that will distinguish between compounds  $\mathbf{Q}$  and  $\mathbf{T}$ , and state the observation made with compound  $\mathbf{T}$ .

**(2)** 

Observation with compound T

(c)	Draw the disp.	layed formula of an	i alkene containing i	tour carbon atoms	•
					(1)

- (d) Three of the compounds belong to the alkane homologous series. All the alkanes in this homologous series have the same general formula.
  - (i) What is the general formula of the alkanes?


(1)

(ii) State two other features of a homologous series.

(2)

```
I ......
```


(e) The displayed formulae below represent isomers.



Explain what isomers are. (2)

(Total for Question 5 = 11 marks)

6 The diagram shows how the electrons are arranged in an atom of oxygen.



Oxygen atoms form both covalent and ionic bonds.

- (a) Water is formed when two atoms of hydrogen combine with one atom of oxygen.
  - (i) Draw a dot and cross diagram of a molecule of water. You need only show the electrons in the outer shells.

(2)

(ii) Explain how the covalent bonds in the water molecule hold the hydrogen and oxygen atoms together.



(i)	Describe, in terms of electrons, what happens when sodium oxide is formed	l in this reaction
		(3)
(ii)	The reaction of sodium to form sodium oxide can be described as oxidation it involves the addition of oxygen.	because
	State one other reason why this reaction can be described as oxidation.	(1)
(c) Exp	plain why water has a much lower melting point than sodium oxide.	(2)
	eacher added sodium oxide to water in a beaker. e equation shows the reaction that occurred.	
	$Na_2O() + H_2O() \rightarrow 2NaOH()$	
	Insert the appropriate state symbols in this equation.	(2)
(i)		
	Some universal indicator was then added to the beaker. A colour change oc State the final colour of the universal indicator and identify the ion responsit the colour change.	
(ii)	State the final colour of the universal indicator and identify the ion responsithe colour change.	
(ii)	State the final colour of the universal indicator and identify the ion responsi	ible for



7	Bromine, chlorine and iodine are elements in Group 7 of the Periodic Table.  (a) (i) Identify which of these elements has	
	(a) (i) Identify which of these elements has	(2)
the	palest colour	
the	e highest melting point	
	(ii) Give the name of another Group 7 element that is a solid at room temperature.	(1)
	(b) When chlorine and hydrogen react together, hydrogen chloride gas forms.	
	Write a chemical equation for this reaction.	(0)
		(2)
	(c) Some hydrogen chloride gas is bubbled into separate samples of water and methylbe A piece of blue litmus paper is dipped into each solution.	nzene.
	(i) State, with a reason, the final colour of the litmus paper in the solution in water.	(2)
	(ii) State, with a reason, the final colour of the litmus paper in the solution in methy	rlbenzene. (2)
	(Total for Question 7 = 9 mas	rks)

8	Some students investigated displacement reactions involving three different metals and
	solutions of their salts. This equation represents one of these reactions:

$$Zn(s) + CuSO_4(aq) \rightarrow ZnSO_4(aq) + Cu(s)$$

This reaction occurs because zinc is more reactive than copper.

When a displacement reaction occurs, there is a temperature rise. The bigger the difference in reactivity between the two metals, the bigger the temperature rise.

(a) What word is used to describe reactions in which there is a temperature rise?

(1)


- (b) The students used this method.
  - Pour some metal salt solution into a beaker, place a thermometer in the beaker and record the temperature
  - Add some of the metal and stir the mixture
  - Record the maximum temperature
  - (i) State **two** variables that the students should keep the same to ensure that the experiment was valid.

/	A	р,
1		,
٧.	-	-

	1	 	 	 	 			 	 	 	 	 	 	 	 	 	 	 		 	 	
١.		 	 	 	 			 	 	 	 	 	 	 	 	 	 	 		 	 	
1	2																					
4	۷	 	 	 	 	• • • • • • •	• • • • • •	 	 	 	 	 • • • • • •	 	 	 	 	 	 	• • • • • • •	 	 	



(ii) The diagrams show the thermometer readings at the start and at the end of one of the experiments.



Record the temperatures and calculate the temperature rise in this experiment.

(3)

Temperature at start

C
Temperature at end

C
Temperature rise

C

(iii) Each experiment was repeated twice. The table shows the average temperatures obtained.

Metal and metal salt used	Average temperature rise in °C
Zn + CuSO <sub>4</sub>	12.2
X + CuSO <sub>4</sub>	8.3
X + ZnSO <sub>4</sub>	0.0
Cu + ZnSO <sub>4</sub>	0.0
Zn + XSO <sub>4</sub>	2.7
Cu + XSO <sub>4</sub>	0.0


Use these results to identify the more reactive metal in each of the following pairs.

(Total for Question 8 =	10 marks)
	(1)
(d) Suggest why the students did not use calcium metal in their experiments.	
	(1)
(c) Write an equation for the reaction with a temperature rise of 2.7 °C.	(1)
Cu and X	
Zn and X	



9 AmmoFert Chemicals is a company that manufactures fertilisers.

The flow chart shows how the company manufactures a fertiliser called AmmoBoost.



- (a) The first step in the process is the conversion of nitrogen gas and hydrogen gas into ammonia.
  - (i) State a raw material used as the source of each gas.

(2)

nitrogen

hydrogen .....

- (ii) Identify the catalyst used in this conversion. (1)
- (iii) State **one** other condition used in this conversion. (1)
- (iv) Only a small percentage of the nitrogen gas and hydrogen gas is converted into ammonia.

Explain how the unreacted gases are separated from the ammonia.



	(b)	The equation	for the	production	of amm	nonia	is
--	-----	--------------	---------	------------	--------	-------	----

$$N_2(g) + 3H_2(g) \rightleftharpoons 2NH_3(g)$$
  $\Delta H = -92 \text{ kJ/mol}$ 

Calculate the maximum mass of ammonia that can be obtained from 56 tonnes of nitrogen. (1 tonne =  $1\,000\,000$  grams)

(3)

- (c) EnAitchThree is another company that manufactures ammonia using the same reaction as AmmoFert but using different conditions.

  EnAitchThree uses a higher temperature and a higher pressure than AmmoFert.
  - (i) Predict the effect on the rate of reaction and on the equilibrium position by changing to the temperature used by EnAitchThree.

(2)

Effect of higher temperature on rate of reaction

Effect of higher temperature on equilibrium position

(ii) Predict the effect on the equilibrium position by changing to the pressure used by EnAitchThree. Justify your prediction.

i) Calculate the percentage by n	nass of oxygen in the compound.	
	,	(1)
ii) Determine the empirical form	nula of the compound.	
		(3)
iii) What is the <b>name</b> of the main	n compound in AmmoBoost?	(1)
		(-)
	(Total for Questi	on 9 = 18 marks)

. , . ,	Explain why iron is malleable.	(2)
		(2)
(ii)	Explain why iron is a good conductor of electricity.	
		(2)
1. \ T	Common Array 100 Array	
	n forms two sulfates. The has the formula $Fe_2(SO_4)_3$	
One		utions
One	has the formula $FeSO_4$ and the other has the formula $Fe_2(SO_4)_3$ addition of sodium hydroxide solution can be used to distinguish between solution	
One The of t	the has the formula $FeSO_4$ and the other has the formula $Fe_2(SO_4)_3$ addition of sodium hydroxide solution can be used to distinguish between solutions sulfates.  State what would be observed in each case.	utions (2)
One The of t	e has the formula $FeSO_4$ and the other has the formula $Fe_2(SO_4)_3$ e addition of sodium hydroxide solution can be used to distinguish between solution established sulfates.	
One The of t	e has the formula FeSO <sub>4</sub> and the other has the formula Fe <sub>2</sub> (SO <sub>4</sub> ) <sub>3</sub> e addition of sodium hydroxide solution can be used to distinguish between solution hese sulfates.  State what would be observed in each case.	
One The of t	the has the formula $FeSO_4$ and the other has the formula $Fe_2(SO_4)_3$ addition of sodium hydroxide solution can be used to distinguish between solutions sulfates.  State what would be observed in each case.	
Ond The of t (i) O <sub>4</sub>	e has the formula FeSO <sub>4</sub> and the other has the formula Fe <sub>2</sub> (SO <sub>4</sub> ) <sub>3</sub> e addition of sodium hydroxide solution can be used to distinguish between solution hese sulfates.  State what would be observed in each case.	
Ond The of t (i) O <sub>4</sub>	e has the formula FeSO <sub>4</sub> and the other has the formula Fe <sub>2</sub> (SO <sub>4</sub> ) <sub>3</sub> e addition of sodium hydroxide solution can be used to distinguish between solution has sulfates.  State what would be observed in each case.  Write a chemical equation for the reaction of iron(II) sulfate (FeSO <sub>4</sub> ) with	
Ond The of t (i) O <sub>4</sub>	e has the formula FeSO <sub>4</sub> and the other has the formula Fe <sub>2</sub> (SO <sub>4</sub> ) <sub>3</sub> e addition of sodium hydroxide solution can be used to distinguish between solution has sulfates.  State what would be observed in each case.  Write a chemical equation for the reaction of iron(II) sulfate (FeSO <sub>4</sub> ) with	(2)



11 Some students investigated the rate of reaction between sodium thiosulfate solution and hydrochloric acid. The equation for the reaction is

$$Na_2S_2O_3(aq) + 2HCl(aq) \rightarrow 2NaCl(aq) + H_2O(l) + S(s) + SO_2(g)$$

The precipitate of sulfur makes the reaction mixture go cloudy.

The students used this method.

- Place a mixture of sodium thiosulfate solution and water in a conical flask
- Add some dilute hydrochloric acid, swirl the mixture and start a timer
- Place the flask over a black cross marked on a piece of paper
- Record the time taken for the cross to disappear when viewed from above

The students used 10 cm<sup>3</sup> of dilute hydrochloric acid in each experiment.

They carried out all the experiments at the same temperature.

They used different volumes of sodium thiosulfate solution and water in each experiment. They were told to keep the total volume of sodium thiosulfate solution and water constant.

The table shows their results.

Student	Volume of sodium thiosulfate solution in cm <sup>3</sup>	Volume of water in cm <sup>3</sup>	Time in s
1	50	0	26.6
2	40	10	55.9
3	35	15	76.4
4	30	20	105.6
5	25	25	149.0
6	20	30	223.5
7	15	40	321.4


(a	) Ex	plain	why	the	results	of	student	7	should	not	be	used
----	------	-------	-----	-----	---------	----	---------	---	--------	-----	----	------

(1)



(b) Plot the results of the six other students on the grid below. Draw a curve of best fit through the points.

(3)



(c) The students used this equation to calculate the rate of each reaction in their investigation.

rate of reaction = 
$$\frac{1000}{\text{time taken}}$$


Calculate the rate of reaction for student 1's experiment.

Give your answer to **one** decimal place.

**(2)** 

Rate = .....

(d) Another group of students used the same method but with different solutions of sodium thiosulfate and hydrochloric acid. They calculated the rate of reaction for each experiment they did. Their results are shown on the following graph.



Rate of reaction in 1000/s

Concentration of sodium thiosulfate solution in g/dm<sup>3</sup>

(i)	Describe the relationship	between rate and	concentration as	shown 1	by the	graph.
-----	---------------------------	------------------	------------------	---------	--------	--------

**(2)** 

(ii) 1	Evnlain	why	increasing	the	concentration	hac	thic	effect	Oη	the	rate

(3)

(Total for Question 11 = 11 marks)

**TOTAL FOR PAPER = 120 MARKS** 

**BLANK PAGE** 



BLANK PAGE

