



UNIVERSITY OF CAMBRIDGE INTERNATIONAL EXAMINATIONS
International General Certificate of Secondary Education

CANDIDATE
NAME

CENTRE
NUMBER

--	--	--	--	--

CANDIDATE
NUMBER

--	--	--	--



CHEMISTRY

0620/31

Paper 3 (Extended)

May/June 2010

1 hour 15 minutes

Candidates answer on the Question Paper.

No Additional Materials are required.

READ THESE INSTRUCTIONS FIRST

Write your Centre number, candidate number and name on all the work you hand in.

Write in dark blue or black pen.

You may use a pencil for any diagrams, graphs or rough working.

Do not use staples, paper clips, highlighters, glue or correction fluid.

DO NOT WRITE IN ANY BARCODES.

Answer **all** questions.

A copy of the Periodic Table is printed on page 16.

At the end of the examination, fasten all your work securely together.

The number of marks is given in brackets [] at the end of each question or part question.

For Examiner's Use	
1	
2	
3	
4	
5	
6	
7	
8	
Total	

This document consists of **13** printed pages and **3** blank pages.



1 Choose an element which fits each of the following descriptions.

(i) It is a yellow solid which burns to form an acidic oxide.

..... [1]

(ii) This element is a black solid which, when heated, forms a purple vapour.

..... [1]

(iii) Most of its soluble salts are blue.

..... [1]

(iv) It has a basic oxide of the type MO which is used to treat acidic soils.

..... [1]

(v) It is an unreactive gas used to fill balloons.

..... [1]

[Total: 5]

2 Ozone is a form of oxygen. Ozone is present in the upper atmosphere and it prevents dangerous solar radiation from reaching the Earth's surface. Some of the chemicals that diffuse into the upper atmosphere decompose ozone. Chemicals that have this effect are methane (CH_4), chloromethane (CH_3Cl) and an oxide of nitrogen (NO_2).

(i) Which of these three chemicals diffuses the most slowly? Give a reason for your choice.

.....

 [2]

(ii) Chloromethane is formed when seaweed decomposes. Name the compounds in the environment from which seaweed might have obtained the following elements:

carbon;

hydrogen;

chlorine. [3]

(iii) How can chloromethane be made from methane?

reagent

condition [2]

3


- (iv) The oxides of nitrogen are atmospheric pollutants. Describe how they are formed.

.....

.....

..... [2]

- (v) Complete the equation for the decomposition of ozone.




[2]

[Total: 11]

For
Examiner's
Use

- 3 A diagram of the apparatus which could be used to investigate the rate of reaction between magnesium and an excess of an acid is drawn below.

For
Examiner's
Use



- (a) The magnesium kept rising to the surface. In one experiment, this was prevented by twisting the magnesium around a piece of copper. In a second experiment, the magnesium was held down by a plastic net fastened to the beaker.


- (i) Suggest a reason why magnesium, which is denser than water, floated to the surface.

.....
..... [1]

- (ii) Iron, zinc and copper have similar densities. Why was copper a better choice than iron or zinc to weigh down the magnesium?

.....
..... [1]

- (b) The only difference in the two experiments was the method used to hold down the magnesium. The results are shown below.



5

For
Examiner's
Use

(i) In which experiment did the magnesium react faster?

..... [1]

(ii) Suggest a reason why the experiment chosen in (i) had the faster rate.

..... [1]

(c) The experiment was repeated using 1.0 mol/dm^3 propanoic acid instead of 1.0 mol/dm^3 hydrochloric acid. Propanoic acid is a weak acid.

(i) How would the graph for propanoic acid **differ** from the graph for hydrochloric acid?

..... [1]

(ii) How would the graph for propanoic acid be the **same** as the graph for hydrochloric acid?

..... [1]

(d) Give **two** factors which would alter the rate of this reaction.

For each factor explain why it alters the rate.

factor

explanation

.....

factor

explanation

..... [4]


[Total: 10]

4 Hydrolysis is used in chemistry to break down complex molecules into simpler ones.

For
Examiner's
Use

(a) Compounds containing the group $\begin{array}{c} \text{O} \\ \parallel \\ \text{—C—} \\ \diagdown \\ \text{O—} \end{array}$ or —COO— are esters.

(i) Give the names and formulae of the two compounds formed when the ester ethyl propanoate is hydrolysed.




name name

formula formula

[4]

(ii) Fats are naturally occurring esters. They can be hydrolysed by boiling with aqueous sodium hydroxide.



What type of compound has the formula $\text{C}_{17}\text{H}_{35}\text{COONa}$ and what is its main use?


type of compound [1]

use [1]

(iii) Name a synthetic polyester.

..... [1]

(b) The structure of a typical protein is drawn below.



(i) What is the name of the polymer linkage?

..... [1]

(ii) Draw the structural formula of a man-made polymer with the same linkage.

[3]

(iii) A protein can be hydrolysed to a mixture of amino acids which are colourless. Individual amino acids can be identified by chromatography. The R_f value of the amino acid glycine is 0.5. Describe how you could show that glycine was present on a chromatogram.

.....

 [3]

[Total: 14]

5 Carbon and silicon are elements in Group IV. Both elements have macromolecular structures.

(a) Diamond and graphite are two forms of the element carbon.

(i) Explain why diamond is a very hard substance.

.....

 [2]

(ii) Give **one** use of diamond.

..... [1]

(iii) Explain why graphite is a soft material.

.....
 [2]

(iv) Give **one** use of graphite.


..... [1]

(b) Two of the oxides of these elements are carbon dioxide, CO_2 , and silicon(IV) oxide, SiO_2 .

(i) Draw a diagram showing the arrangement of the valency electrons in one molecule of the covalent compound carbon dioxide.
 Use x to represent an electron from a carbon atom.
 Use o to represent an electron from an oxygen atom.

[3]

(ii) A section of the macromolecular structure of silicon(IV) oxide is given below.



Use this diagram to explain why the formula is SiO_2 not SiO_4 .


.....
 [2]

(iii) Predict **two** differences in the physical properties of these two oxides.


.....
 [2]

[Total: 13]

- 6 Iodine reacts with chlorine to form dark brown iodine monochloride.



This reacts with more chlorine to give yellow iodine trichloride.
There is an equilibrium between these iodine chlorides.



For
Examiner's
Use

- (a) Explain what is meant by *equilibrium*.

.....
.....
..... [2]

- (b) When the equilibrium mixture is heated it becomes a darker brown colour.
Is the reverse reaction endothermic or exothermic? Give a reason for your choice.

.....
.....
..... [2]

- (c) The pressure on the equilibrium mixture is decreased.

- (i) How would this affect the position of equilibrium and why?

It would move to the [1]

reason


..... [1]

- (ii) Describe what you would observe.

.....
..... [1]

[Total: 7]


- 7 Titanium is a transition element. It is isolated by the following reactions.



- (a) Why is it usually necessary to include a number in the name of the compounds of transition elements?

..... [1]

- (b) Titanium(IV) chloride is made by heating the oxide with coke and chlorine.



Explain why the presence of coke ensures the maximum yield of the metal chloride.

.....

 [2]

- (c) Explain why the change, titanium(IV) chloride to titanium, is reduction.

.....
 [1]

- (d) Complete the table which shows some of the properties of titanium and its uses. The first line has been completed as an example.

property	related use
soluble in molten steel	making steel titanium alloys
.....	making aircraft and space vehicles
resistant to corrosion, especially in sea water

[2]

(e) The titanium ore contains 36.8% iron, 31.6% titanium and the remainder is oxygen.

(i) Determine the percentage of oxygen in this titanium compound.

percentage of oxygen = % [1]

(ii) Calculate the number of moles of atoms for each element.

The number of moles of Fe is shown as an example.

number of moles of Fe = $36.8/56 = 0.66$

number of moles of Ti =

number of moles of O = [1]

(iii) What is the simplest ratio for the moles of atoms?

Fe	:	Ti	:	O
.....	

[1]

(iv) What is the formula of this titanium compound?

..... [1]

[Total: 10]

8 Methanoic acid is the first member of the homologous series of carboxylic acids.


(a) Give **two** general characteristics of a homologous series.

.....

 [2]

(b) In some areas when water is boiled, the inside of kettles become coated with a layer of calcium carbonate. This can be removed by adding methanoic acid.

(i) Complete the equation.



(ii) Methanoic acid reacts with most metals above hydrogen in the reactivity series. Complete the word equation.

zinc + methanoic acid \rightarrow + [2]

(iii) Aluminium is also above hydrogen in the reactivity series. Why does methanoic acid not react with an aluminium kettle?

.....
 [1]

(c) Give the name, molecular formula and empirical formula of the fourth acid in this series.

name [1]

molecular formula [1]

empirical formula [1]

[Total: 10]

DATA SHEET
The Periodic Table of the Elements

Group		I	II	III	IV	V	VI	VII	0														
7	9	1								4													
Li Lithium 3	Be Beryllium 4	H Hydrogen 1								He Helium 2													
23	24								19	20													
Na Sodium 11	Mg Magnesium 12								F Fluorine 9	Ne Neon 10													
39	40								16	17													
K Potassium 19	Ca Calcium 20								O Oxygen 8	Cl Chlorine 17													
85	88								14	15													
Rb Rubidium 37	Sr Strontium 38								N Nitrogen 7	P Phosphorus 15													
133	137								28	31													
Cs Caesium 55	Ba Barium 56								Al Aluminium 13	Si Silicon 14													
226	227								70	73													
Fr Francium 87	Ra Radium 88								Ga Gallium 31	Ge Germanium 32													
		59	59	59	59	59	59	59	80														
		Sc Scandium 21	Ti Titanium 22	V Vanadium 23	Cr Chromium 24	Mn Manganese 25	Fe Iron 26	Co Cobalt 27	Ni Nickel 28	Cu Copper 29	Zn Zinc 30	Se Selenium 34	Br Bromine 35	Kr Krypton 36									
		89	91	93	96	101	101	103	106	108	112	128	127	131									
		Y Yttrium 39	Zr Zirconium 40	Nb Niobium 41	Mo Molybdenum 42	Tc Technetium 43	Ru Ruthenium 44	Rh Rhodium 45	Pd Palladium 46	Ag Silver 47	Cd Cadmium 48	Te Tellurium 52	I Iodine 53	Xe Xenon 54									
		139	178	181	184	186	190	192	195	197	201	207	209	209									
		La Lanthanum 57	Hf Hafnium 72	Ta Tantalum 73	W Tungsten 74	Re Rhenium 75	Os Osmium 76	Ir Iridium 77	Pt Platinum 78	Au Gold 79	Hg Mercury 80	Pb Lead 82	Bi Bismuth 83	Rn Radon 86									
		226								84	85	86											
		Ac Actinium 89								Po Polonium 84	At Astatine 85	Rn Radon 86											
												165	166	167	169	173	175						
												Ho Holmium 67	Er Erbium 68	Tm Thulium 69	Yb Ytterbium 70	Lu Lutetium 71	175						
												162	162	162	162	162	162	162					
												Dy Dysprosium 66	Ho Holmium 67	Er Erbium 68	Tm Thulium 69	Yb Ytterbium 70	Lu Lutetium 71	175					
												159	159	159	159	159	159	159					
												Tb Terbium 65	Dy Dysprosium 66	Ho Holmium 67	Er Erbium 68	Tm Thulium 69	Yb Ytterbium 70	Lu Lutetium 71	175				
												157	157	157	157	157	157	157					
												Gd Gadolinium 64	Tb Terbium 65	Dy Dysprosium 66	Ho Holmium 67	Er Erbium 68	Tm Thulium 69	Yb Ytterbium 70	Lu Lutetium 71	175			
												152	152	152	152	152	152	152					
												Eu Europium 63	Gd Gadolinium 64	Tb Terbium 65	Dy Dysprosium 66	Ho Holmium 67	Er Erbium 68	Tm Thulium 69	Yb Ytterbium 70	Lu Lutetium 71	175		
												150	150	150	150	150	150	150					
												Sm Samarium 62	Eu Europium 63	Gd Gadolinium 64	Tb Terbium 65	Dy Dysprosium 66	Ho Holmium 67	Er Erbium 68	Tm Thulium 69	Yb Ytterbium 70	Lu Lutetium 71	175	
												144	144	144	144	144	144	144					
												Pm Promethium 61	Sm Samarium 62	Eu Europium 63	Gd Gadolinium 64	Tb Terbium 65	Dy Dysprosium 66	Ho Holmium 67	Er Erbium 68	Tm Thulium 69	Yb Ytterbium 70	Lu Lutetium 71	175
												238	238	238	238	238	238	238					
												Np Neptunium 93	Pu Plutonium 94	Am Americium 95	Cm Curium 96	Bk Berkelium 97	Cf Californium 98	Es Einsteinium 99	Fm Fermium 100	Md Mendelevium 101	No Nobelium 102	Lr Lawrencium 103	175
		140	141	144	152	157	159	162	165	167	169	173	175	175									
		Ce Cerium 58	Pr Praseodymium 59	Nd Neodymium 60	Pm Promethium 61	Sm Samarium 62	Eu Europium 63	Gd Gadolinium 64	Tb Terbium 65	Dy Dysprosium 66	Ho Holmium 67	Er Erbium 68	Tm Thulium 69	Yb Ytterbium 70	Lu Lutetium 71	175							
		232	232	238	238	238	238	238	238	238	238	238	238	238	238	238							
		Th Thorium 90	Pa Protactinium 91	U Uranium 92	Np Neptunium 93	Pu Plutonium 94	Am Americium 95	Cm Curium 96	Bk Berkelium 97	Cf Californium 98	Es Einsteinium 99	Fm Fermium 100	Md Mendelevium 101	No Nobelium 102	Lr Lawrencium 103	175							

The volume of one mole of any gas is 24 dm³ at room temperature and pressure (r.t.p.).

a	X	b
Key		
a = relative atomic mass	X = atomic symbol	b = proton (atomic) number

*58-71 Lanthanoid series
†90-103 Actinoid series

Permission to reproduce items where third-party owned material protected by copyright is included has been sought and cleared where possible. Every reasonable effort has been made by the publisher (UCLES) to trace copyright holders, but if any items requiring clearance have unwittingly been included the publisher will be pleased to make amends at the earliest possible opportunity.

University of Cambridge International Examinations is part of the Cambridge Assessment Group. Cambridge Assessment is the brand name of University of Cambridge Local Examinations Syndicate (UCLES), which is itself a department of the University of Cambridge.