



# Cambridge IGCSE™

---

**PHYSICS****0625/42**

Paper 4 Theory (Extended)

**March 2021**

MARK SCHEME

Maximum Mark: 80

---

<b>Published</b>
------------------

This mark scheme is published as an aid to teachers and candidates, to indicate the requirements of the examination. It shows the basis on which Examiners were instructed to award marks. It does not indicate the details of the discussions that took place at an Examiners' meeting before marking began, which would have considered the acceptability of alternative answers.

Mark schemes should be read in conjunction with the question paper and the Principal Examiner Report for Teachers.

Cambridge International will not enter into discussions about these mark schemes.

Cambridge International is publishing the mark schemes for the March 2021 series for most Cambridge IGCSE™, Cambridge International A and AS Level components and some Cambridge O Level components.

---

This document consists of **10** printed pages.

**PUBLISHED****Generic Marking Principles**

These general marking principles must be applied by all examiners when marking candidate answers. They should be applied alongside the specific content of the mark scheme or generic level descriptors for a question. Each question paper and mark scheme will also comply with these marking principles.

**GENERIC MARKING PRINCIPLE 1:**

Marks must be awarded in line with:

- the specific content of the mark scheme or the generic level descriptors for the question
- the specific skills defined in the mark scheme or in the generic level descriptors for the question
- the standard of response required by a candidate as exemplified by the standardisation scripts.

**GENERIC MARKING PRINCIPLE 2:**

Marks awarded are always **whole marks** (not half marks, or other fractions).

**GENERIC MARKING PRINCIPLE 3:**

Marks must be awarded **positively**:

- marks are awarded for correct/valid answers, as defined in the mark scheme. However, credit is given for valid answers which go beyond the scope of the syllabus and mark scheme, referring to your Team Leader as appropriate
- marks are awarded when candidates clearly demonstrate what they know and can do
- marks are not deducted for errors
- marks are not deducted for omissions
- answers should only be judged on the quality of spelling, punctuation and grammar when these features are specifically assessed by the question as indicated by the mark scheme. The meaning, however, should be unambiguous.

**GENERIC MARKING PRINCIPLE 4:**

Rules must be applied consistently, e.g. in situations where candidates have not followed instructions or in the application of generic level descriptors.

**GENERIC MARKING PRINCIPLE 5:**

Marks should be awarded using the full range of marks defined in the mark scheme for the question (however; the use of the full mark range may be limited according to the quality of the candidate responses seen).

**GENERIC MARKING PRINCIPLE 6:**

Marks awarded are based solely on the requirements as defined in the mark scheme. Marks should not be awarded with grade thresholds or grade descriptors in mind.

**Science-Specific Marking Principles**

- 1 Examiners should consider the context and scientific use of any keywords when awarding marks. Although keywords may be present, marks should not be awarded if the keywords are used incorrectly.
- 2 The examiner should not choose between contradictory statements given in the same question part, and credit should not be awarded for any correct statement that is contradicted within the same question part. Wrong science that is irrelevant to the question should be ignored.
- 3 Although spellings do not have to be correct, spellings of syllabus terms must allow for clear and unambiguous separation from other syllabus terms with which they may be confused (e.g. ethane / ethene, glucagon / glycogen, refraction / reflection).
- 4 The error carried forward (ecf) principle should be applied, where appropriate. If an incorrect answer is subsequently used in a scientifically correct way, the candidate should be awarded these subsequent marking points. Further guidance will be included in the mark scheme where necessary and any exceptions to this general principle will be noted.
- 5 'List rule' guidance  
For questions that require **n** responses (e.g. State **two** reasons ...):
  - The response should be read as continuous prose, even when numbered answer spaces are provided.
  - Any response marked *ignore* in the mark scheme should not count towards **n**.
  - Incorrect responses should not be awarded credit but will still count towards **n**.
  - Read the entire response to check for any responses that contradict those that would otherwise be credited. Credit should **not** be awarded for any responses that are contradicted within the rest of the response. Where two responses contradict one another, this should be treated as a single incorrect response.
  - Non-contradictory responses after the first **n** responses may be ignored even if they include incorrect science.

**6** Calculation specific guidance

Correct answers to calculations should be given full credit even if there is no working or incorrect working, **unless** the question states 'show your working'.

For questions in which the number of significant figures required is not stated, credit should be awarded for correct answers when rounded by the examiner to the number of significant figures given in the mark scheme. This may not apply to measured values.

For answers given in standard form (e.g.  $a \times 10^n$ ) in which the convention of restricting the value of the coefficient ( $a$ ) to a value between 1 and 10 is not followed, credit may still be awarded if the answer can be converted to the answer given in the mark scheme.

Unless a separate mark is given for a unit, a missing or incorrect unit will normally mean that the final calculation mark is not awarded. Exceptions to this general principle will be noted in the mark scheme.

**7** Guidance for chemical equations

Multiples / fractions of coefficients used in chemical equations are acceptable unless stated otherwise in the mark scheme.

State symbols given in an equation should be ignored unless asked for in the question or stated otherwise in the mark scheme.

Question	Answer	Marks
1(a)	78 N	<b>A3</b>
	( $m=$ ) $\rho V$ OR $\rho = m / V$ in any form	C1
	$W = mg$	C1
1(b)	$4.5 \times 10^3$ N	<b>A3</b>
	( $F=$ ) $(\Delta)PA$ OR $P = F / A$ in any form	C1
	( $\Delta P = 1.3 \times 10^5 - 1.0 \times 10^5 =$ ) $3 \times 10^4$	C1
	outwards	<b>B1</b>
1(c)	( $\rho=$ ) $800 \text{ kg / m}^3$	<b>A3</b>
	( $\rho=$ ) $P / gh$ OR $P = \rho gh$ in any form	C1
	( $\rho=$ ) $9.6 \times 10^4 / (10 \times 12)$	C1

Question	Answer	Marks
2(a)(i)	(moment of a force) is the turning effect (about a point / pivot)	<b>B1</b>
2(a)(ii)	$2.8 \times 10^6$ N m	<b>A2</b>
	(moment = ) $Fd$ in any form	C1
2(b)(i)	scalar / speed has magnitude <u>only</u> OR scalar / speed has no direction	<b>B1</b>
	vector / velocity has magnitude and direction	<b>B1</b>
2(b)(ii)	any scalar quantity	<b>B1</b>
	any vector quantity	<b>B1</b>

Question	Answer	Marks
2(c)	correct triangle or parallelogram drawn	<b>B1</b>
	resultant force (including correct arrow)	<b>B1</b>
	scale 1 cm = 4 N or 1 cm = 5 N	<b>B1</b>
	40–47 N <b>AND</b> 33°–40° (anticlockwise from 20 N)	<b>B1</b>

Question	Answer	Marks
3(a)	renewable / yes	<b>B1</b>
	crops can be regrown (to replace resource) / waste materials don't run out	<b>B1</b>
3(b)	water will cool (too much) / thermal energy lost (during transfer)	<b>B1</b>
	lag/insulate (pipes) OR transport in a poor conductor of thermal energy	<b>B1</b>
3(c)	any two from: <ul style="list-style-type: none"> <li>• air pollution / harmful gases / acid rain</li> <li>• CO<sub>2</sub> / greenhouse gases / contribution to global warming</li> <li>• not renewable</li> <li>• damage from mining / drilling or any valid environmental consequence of transport of coal</li> </ul>	<b>B2</b>

Question	Answer	Marks
4(a)	molecules strike walls	<b>B1</b>
	momentum (of molecules) changes / momentum = mass × velocity	<b>B1</b>
	force = rate of change of momentum	<b>B1</b>
	pressure = (sum of) force(s) / area / pressure = rate of change of momentum / area	<b>B1</b>

Question	Answer	Marks
4(b)(i)	$(p_2=) p_1 V_1 / V_2$	<b>A2</b>
	$p_1 V_1 = p_2 V_2$	C1
4(b)(ii)	<u>greater</u>	<b>B1</b>
	molecules move <u>faster</u> / have <u>greater</u> KE / molecules have <u>greater</u> momentum	<b>B1</b>
	(leads to) <u>more</u> frequent / <u>harder</u> collisions (with walls) / great rate of change of momentum	<b>B1</b>

Question	Answer	Marks
5(a)	echo	<b>B1</b>
5(b)	$(\lambda=) 7.5 \times 10^{-4} \text{ m}$	<b>A3</b>
	$(\lambda=) v / f$ OR $v = f\lambda$ in any form	C1
	$(\lambda=) 1.5 \times 10^3 / 2 \times 10^6$	C1
5(c)(i),(ii)	labelled wavelength of incident wave	<b>B1</b>
	3 part circles to the left of the barrier and centred to right of the barrier	<b>B1</b>
	wavelengths of reflected and incident waves same	<b>B1</b>

Question	Answer	Marks
6(a)	Any two correct rays from <ul style="list-style-type: none"> <li>• from O through optical centre (and beyond)</li> <li>• from O parallel to principal axis to centre line of lens then through <math>F_1</math></li> <li>• from <math>F_2</math> through O to centreline of lens then parallel to principal axis</li> </ul>	<b>M2</b>
	rays traced back to intersect <b>AND</b> 2.4 – 3.6 cm	<b>A1</b>
6(b)	magnified	<b>B1</b>
	same way up as object	<b>B1</b>
	virtual	<b>B1</b>
6(c)	one ray from each prism refracted towards principal axis	<b>B1</b>
	(rays) converge to the right of original convergence on the principal axis	<b>B1</b>

Question	Answer	Marks
7(a)	no cutting of (magnetic) flux / <u>magnetic</u> field	<b>B1</b>
7(b)	to the top of the page / RH box	<b>B1</b>
	current, motion and (magnetic) field mutually at right angles	<b>B1</b>
	(magnetic) field from left to right	<b>B1</b>
7(c)(i)	opposite current (direction) / opposite deflection (on ammeter)	<b>B1</b>
7(c)(ii)	greater current / deflection	<b>B1</b>



Question	Answer	Marks
8(a)	<u>energy</u> supplied by a source in driving charge around a complete circuit / <u>energy</u> needed to drive unit charge / 1 coulomb round circuit	<b>B1</b>
8(b)(i)	$(P=) 90 \text{ W}$	<b>A3</b>
	$(P=) VI$ in any form	C1
	$(V/R \text{ OR } I =) 2$	C1
8(b)(ii)	(p.d. =) 15 V	<b>A2</b>
	(p.d. =) 60–45	C1
8(b)(iii)	$(I = 15 / 10 =) 1.5 \text{ A}$	<b>A2</b>
	$(I =) V / R \text{ OR } V = IR$ in any form	<b>C1</b>

Question	Answer	Marks															
9(a)	<table border="0"> <tr> <td>I/P</td> <td>I/P</td> <td>O/P</td> </tr> <tr> <td>0</td> <td>0</td> <td></td> </tr> <tr> <td>0</td> <td>1</td> <td></td> </tr> <tr> <td>1</td> <td>0</td> <td></td> </tr> <tr> <td>1</td> <td>1</td> <td></td> </tr> </table>	I/P	I/P	O/P	0	0		0	1		1	0		1	1		<b>B1</b>
	I/P	I/P	O/P														
0	0																
0	1																
1	0																
1	1																
	<table border="0"> <tr> <td>I/P</td> <td>I/P</td> <td>O/P</td> </tr> <tr> <td>0</td> <td>0</td> <td><b>0</b></td> </tr> <tr> <td>0</td> <td>1</td> <td><b>1</b></td> </tr> <tr> <td>1</td> <td>0</td> <td><b>1</b></td> </tr> <tr> <td>1</td> <td>1</td> <td><b>1</b></td> </tr> </table>	I/P	I/P	O/P	0	0	<b>0</b>	0	1	<b>1</b>	1	0	<b>1</b>	1	1	<b>1</b>	<b>B1</b>
I/P	I/P	O/P															
0	0	<b>0</b>															
0	1	<b>1</b>															
1	0	<b>1</b>															
1	1	<b>1</b>															
9(b)	two inputs to curved face, sharp end with small circle and one output	<b>B1</b>															

Question	Answer	Marks
9(c)(i)	1 0 1 0	
9(c)(ii)	AND	<b>B1</b>
	input 1 and 1 gives output 1	<b>B1</b>
	any 0 input gives 0 output	<b>B1</b>

Question	Answer	Marks
10(a)	2	<b>B1</b>
	4	<b>B1</b>
	+2	<b>B1</b>
10(b)	${}_{38}^{90}\text{Sr} \rightarrow {}_{39}^{90}\text{Y} + {}_{-1}^0\beta$	
	nucleon numbers 90 on both sides of equation	<b>B1</b>
	Sr and proton number 38 on left AND Y and proton number 39 on right	<b>B1</b>
	${}_{-1}^0\beta$ (to right of arrow)	<b>B1</b>
10(c)	(original mass = $4 / 9.2 =$ ) 37 mg	<b>A2</b>
	2 half-lives	<b>C1</b>