



Cambridge International Examinations

Cambridge International General Certificate of Secondary Education

CANDIDATE NAME						
CENTRE NUMBER				CANDIDATE NUMBER		

CHEMISTRY 0620/51

Paper 5 Practical Test October/November 2015

1 hour 15 minutes

Candidates answer on the Question Paper.

Additional Materials: As listed in the Confidential Instructions

READ THESE INSTRUCTIONS FIRST

Write your Centre number, candidate number and name on all the work you hand in.

Write in dark blue or black pen.

You may use an HB pencil for any diagrams or graphs.

Do not use staples, paper clips, glue or correction fluid.

DO NOT WRITE IN ANY BARCODES.

Answer all questions.

Electronic calculators may be used.

You may lose marks if you do not show your working or if you do not use appropriate units.

Practical notes are provided on page 8.

At the end of the examination, fasten all your work securely together.

The number of marks is given in brackets [] at the end of each question or part question.

For Examiner's Use		
Total		

The syllabus is approved for use in England, Wales and Northern Ireland as a Cambridge International Level 1/Level 2 Certificate.

This document consists of 7 printed pages and 1 blank page.



1 You are going to investigate the rate of reaction when magnesium ribbon reacts with four solutions of dilute sulfuric acid, of different concentrations, labelled solutions **A**, **B**, **C** and **D**. The sulfuric acid is in excess in all experiments.

Read all the instructions below carefully before starting the experiments.

Instructions

You are going to carry out five experiments.

(a) Experiment 1

Using a measuring cylinder, pour 30 cm³ of dilute sulfuric acid **A** into the beaker.

Start the timer and add the 4cm length of magnesium to the sulfuric acid in the beaker. Stir the mixture constantly.

Measure the time taken for all of the magnesium to react and disappear.

Record the time in the table below.

Rinse out the beaker with distilled water.

(b) Experiment 2

Repeat Experiment 1, using the solution **B** of sulfuric acid. Record the time in the table.

(c) Experiment 3

Repeat Experiment 1, using the solution **C** of sulfuric acid. Record the time in the table.

(d) Experiment 4

Repeat Experiment 1, using the solution **D** of sulfuric acid. Record the time in the table.


Experiment number	solution of sulfuric acid	concentration of sulfuric acid in mol/dm³	time for magnesium to completely disappear in seconds
1	Α	1.0	
2	В	0.8	
3	С	0.6	
4	D	0.5	


[4]

(e) Experiment 5

Pour about 3 cm ³ of the solution	C of sulfuric acid in	nto a test-tube and	record its temperature.
Add a 4 cm length of magnesiun	٦.		

Record all of your observations and test the gas given off with a splint.





[3]

(g) (i) From your graph, deduce the concentration of the sulfuric acid if the time for the reaction was 80 s.

Show clearly on the graph how you worked out your answer.

..... mol/dm³ [2]

4

	(ii)	From your graph , deduce how long the reaction would take if a solution of sulfuric acid of concentration 1.2 mol/dm³ was used. Show clearly on the graph how you worked out your answer.
		s [2]
(h)) Wh	ny was the same amount of magnesium used in Experiments 1, 2, 3 and 4?
	••••	[1]
(i)	Us	e your observations in Experiment 5 to answer these questions.
	(i)	What type of chemical reaction occurs when magnesium reacts with sulfuric acid?
		[1]
	(ii)	Identify the gas given off.
		[1]
	(iii)	Suggest the effect on the temperature change if this experiment was repeated using 2 cm of magnesium ribbon.
		[1]
(j)	ma	ggest a different method which could be used to investigate the rate of the reaction between agnesium and sulfuric acid. State the difference in the apparatus used and measurements to taken.
	apı	paratus
	me	easurements
		[3]
		[Total: 20]

You are provided with two metal salt solutions, **X** and **Y**.

Carry out the following tests on **X** and **Y**, recording all of your observations in the tables.

Conclusions must **not** be written in the tables.

	tests	observations
test	s on solution X	
(a)	Describe the appearance of solution X .	[1]
1	ide the solution into four equal portions in arate test-tubes.	
(b)	To the first portion of solution add about 1 cm ³ of dilute nitric acid followed by aqueous silver nitrate.	[2]
(c)	To the second portion of solution add about 1 cm³ of aqueous sodium hydroxide and shake the mixture and leave to stand.	[1]
	Note how the mixture changes after five minutes.	[1]
(d)	To the third portion of solution add excess aqueous ammonia.	[1]
(e)	To the fourth portion of solution add a few drops of dilute sulfuric acid followed by about 1 cm ³ of aqueous potassium manganate(VII).	[1]
	Now add aqueous sodium hydroxide to the mixture.	[2]

tests	observations
tests on solution Y	
Divide the solution into three equal poseparate test-tubes.	ortions in
(f) To the first portion of the solution at 1 cm ³ of dilute hydrochloric acid.	dd about[1]
(g) Tip the second portion of the solution a boiling tube. Add about 1 cm³ of sodium hydroxide and shake the radd one small piece of aluminiuthe mixture and warm gently. Test given off with red litmus paper.	aqueous nixture. m foil to
(h) To the third portion of the soluraqueous potassium iodide.	tion add[2]
i) What conclusions can you draw ab	out solution X ?
	[3]
(j) What conclusions can you draw ab	out solution Y ?
	[2]
	[Total: 20]

7

BLANK PAGE

8

NOTES FOR USE IN QUALITATIVE ANALYSIS

Test for anions

anion	test	test result
carbonate (CO ₃ ²⁻)	add dilute acid	effervescence, carbon dioxide produced
chloride (C <i>l</i> ⁻) [in solution]	acidify with dilute nitric acid, then add aqueous silver nitrate	white ppt.
iodide (I ⁻) [in solution]	acidify with dilute nitric acid, then add aqueous silver nitrate	yellow ppt.
nitrate (NO ₃ ⁻) [in solution]	add aqueous sodium hydroxide then aluminium foil; warm carefully	ammonia produced
sulfate (SO ₄ ²⁻) [in solution]	acidify with dilute nitric acid, then aqueous barium nitrate	white ppt.

Test for aqueous cations

cation	effect of aqueous sodium hydroxide	effect of aqueous ammonia
aluminium (Al ³⁺)	white ppt., soluble in excess giving a colourless solution	white ppt., insoluble in excess
ammonium (NH ₄ +)	ammonia produced on warming	_
calcium (Ca ²⁺)	white ppt., insoluble in excess	no ppt., or very slight white ppt.
copper (Cu ²⁺)	light blue ppt., insoluble in excess	light blue ppt., soluble in excess giving a dark blue solution
iron(II) (Fe ²⁺)	green ppt., insoluble in excess	green ppt., insoluble in excess
iron(III) (Fe ³⁺)	red-brown ppt., insoluble in excess	red-brown ppt., insoluble in excess
zinc (Zn ²⁺)	white ppt., soluble in excess giving a colourless solution	white ppt., soluble in excess giving a colourless solution

Test for gases

gas	test and test results	
ammonia (NH ₃)	turns damp red litmus paper blue	
carbon dioxide (CO ₂)	turns limewater milky	
chlorine (Cl ₂)	bleaches damp litmus paper	
hydrogen (H ₂)	'pops' with a lighted splint	
oxygen (O ₂)	relights a glowing splint	

Permission to reproduce items where third-party owned material protected by copyright is included has been sought and cleared where possible. Every reasonable effort has been made by the publisher (UCLES) to trace copyright holders, but if any items requiring clearance have unwittingly been included, the publisher will be pleased to make amends at the earliest possible opportunity.

To avoid the issue of disclosure of answer-related information to candidates, all copyright acknowledgements are reproduced online in the Cambridge International Examinations Copyright Acknowledgements Booklet. This is produced for each series of examinations and is freely available to download at www.cie.org.uk after the live examination series.

Cambridge International Examinations is part of the Cambridge Assessment Group. Cambridge Assessment is the brand name of University of Cambridge Local Examinations Syndicate (UCLES), which is itself a department of the University of Cambridge.