

Write your name here

Surname					Other names				
Centre Number					Candidate Number				
Pearson Edexcel International GCSE (9 - 1)					<input type="text"/> <input type="text"/> <input type="text"/> <input type="text"/> <input type="text"/>				
<h1>Chemistry</h1> <h2>Paper 2</h2>									
Sample Assessment Materials for first teaching September 2017 Time: 1 hour 15 minutes					Paper Reference 4CH1/2C				
You must have: Calculator, ruler					Total Marks <input type="text"/>				

Instructions

- Use **black** ink or ball-point pen.
- **Fill in the boxes** at the top of this page with your name, centre number and candidate number.
- Answer **all** questions.
- Answer the questions in the spaces provided
– *there may be more space than you need.*
- Calculators may be used.
- Some questions must be answered with a cross in a box ☒. If you change your mind about an answer, put a line through the box ☒ and then mark your new answer with a cross ☒.

Information

- The total mark for this paper is 70.
- The marks for **each** question are shown in brackets
– *use this as a guide as to how much time to spend on each question.*

Advice

- Read each question carefully before you start to answer it.
- Try to answer every question.
- Check your answers if you have time at the end.

Turn over ►

S52916A

©2016 Pearson Education Ltd.



PEARSON

The Periodic Table of the Elements

1	2	3	4	5	6	7	0																																																																										
7 Li lithium 3	9 Be beryllium 4	11 Na sodium 11	12 C carbon 6	13 Al aluminium 13	14 N nitrogen 7	15 P phosphorus 15	16 O oxygen 8	17 Cl chlorine 17	18 Ar argon 18	19 F fluorine 9	20 Ne neon 10	21 K potassium 19	22 Ca calcium 20	23 V vanadium 23	24 Cr chromium 24	25 Mn manganese 25	26 Fe iron 26	27 Co cobalt 27	28 Ni nickel 28	29 Cu copper 29	30 Zn zinc 30	31 Ga gallium 31	32 Ge germanium 32	33 As arsenic 33	34 Se selenium 34	35 Br bromine 35	36 Kr krypton 36	37 Rb rubidium 37	38 Sr strontium 38	39 Y yttrium 39	40 Zr zirconium 40	41 Nb niobium 41	42 Mo molybdenum 42	43 Tc technetium [98]	44 Ru ruthenium 44	45 Rh rhodium 45	46 Pd palladium 46	47 Ag silver 47	48 Cd cadmium 48	49 In indium 49	50 Sn tin 50	51 Sb antimony 51	52 Te tellurium 52	53 I iodine 53	54 Xe xenon 54	55 Cs caesium 55	56 Ba barium 56	57 La* lanthanum 57	72 Hf hafnium 72	73 Ta tantalum 73	74 W tungsten 74	75 Re rhenium 75	76 Os osmium 76	77 Ir iridium 77	78 Pt platinum 78	79 Au gold 79	80 Hg mercury 80	81 Tl thallium 81	82 Pb lead 82	83 Bi bismuth 83	84 Po polonium 84	85 At astatine 85	86 Rn radon 86	87 Fr francium 87	88 Ra radium 88	89 Ac* actinium 89	104 Rf rutherfordium [261]	105 Db dubnium [262]	106 Sg seaborgium [266]	107 Bh bohrium [264]	108 Hs hassium [277]	109 Mt meitnerium [268]	110 Ds darmstadtium [271]	111 Rg roentgenium [272]	112 Cn copernicium [285]	113 Nh nihonium [286]	114 Fl flerovium [289]	115 Mc moscovium [288]	116 Lv livermorium [293]	117 Ts tennessine [294]	118 Og oganesson [294]
												<p style="text-align: center;">Key</p> <table border="1" style="margin-left: auto; margin-right: auto;"> <tr> <td style="padding: 5px;">relative atomic mass</td> </tr> <tr> <td style="padding: 5px;">atomic symbol <small>name</small></td> </tr> <tr> <td style="padding: 5px;">atomic (proton) number</td> </tr> </table>												relative atomic mass	atomic symbol <small>name</small>	atomic (proton) number																																																							
relative atomic mass																																																																																	
atomic symbol <small>name</small>																																																																																	
atomic (proton) number																																																																																	
												<p>Elements with atomic numbers 112-116 have been reported but not fully authenticated</p>																																																																					

* The lanthanoids (atomic numbers 58-71) and the actinoids (atomic numbers 90-103) have been omitted.

The relative atomic masses of copper and chlorine have not been rounded to the nearest whole number.

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

BLANK PAGE

Answer ALL questions. Write your answers in the spaces provided.

- 1 The table shows some properties of four substances, P, Q, R and S.

Substance	Melting point /°C	Boiling point /°C	Conducts electricity when	
			solid	liquid
P	3410	5930	yes	yes
Q	734	1435	no	yes
R	-95	69	no	no
S	2507	3900	no	no

Use the information in the table to answer the following questions.
You may use each letter once, more than once or not at all.

Choose a substance that:

- (a) is a solid at 3000 °C.

(1)

- A substance P
- B substance Q
- C substance R
- D substance S

- (b) is a liquid at 25 °C.

(1)

- A substance P
- B substance Q
- C substance R
- D substance S

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

(c) is an ionic compound.

(1)

- A substance P
- B substance Q
- C substance R
- D substance S

(d) is a metal.

(1)

- A substance P
- B substance Q
- C substance R
- D substance S

(Total for Question 1 = 4 marks)

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

2 This is a method used to measure the solubility of a solid in water:

- add an excess of solid to some water in a boiling tube and stir
- measure the temperature of the saturated solution formed
- weigh an empty evaporating basin
- pour some of the saturated solution into the evaporating basin
- weigh the basin and contents
- heat the evaporating basin to remove all of the water
- weigh the evaporating basin and remaining solid.

(a) The table shows the results of an experiment using this method.

mass of evaporating basin / g	89.6
mass of evaporating basin + saturated solution / g	115.8
mass of evaporating basin + solid / g	94.9

Calculate the mass of solid obtained and the mass of water removed.

(2)

mass of solid = g

mass of water = g

(b) In another experiment, at a different temperature, the mass of solid obtained is 10.5 g and the mass of water removed is 16.8 g.

Calculate the solubility of the solid, in g per 100 g of water, at this temperature.

(2)

solubility = g per 100 g of water

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

- (c) If the evaporating basin is heated too strongly some of the solid decomposes to form a gas.

Explain how this strong heating would affect the value of the calculated solubility of the solid.

(3)

.....

.....

.....

.....

.....

.....

.....

(Total for Question 2 = 7 marks)

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

3 Astatine, bromine, chlorine, fluorine and iodine are all halogens.
They are found in Group 7 of the Periodic Table.

(a) Predict which halogen has the lightest colour.

(1)

(b) Name a halogen that is a solid at room temperature.

(1)


(c) Bromine can be obtained from the bromide ions in sea water.

Chlorine is bubbled into sea water.

The chlorine oxidises the bromide ions to bromine atoms.

The bromine atoms then form bromine molecules.

(i) Complete the equation to show how bromine **atoms** are formed from bromide ions.



(1)

(ii) State why this reaction is described as the oxidation of bromide ions.

(1)

(iii) Write an equation to show how bromine atoms form bromine molecules.

(1)

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

(d) Boron and fluorine form a covalent compound that has the molecular formula BF_3

Draw a dot-and-cross diagram to show the arrangement of the outer electrons in a molecule of BF_3

Use crosses (X) to represent the outer electrons of boron. Use dots (•) to represent the outer electrons of fluorine.

(2)

(Total for Question 3 = 7 marks)

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA



BLANK PAGE

5 Zinc metal is obtained from sulfide ores.



The most common ore of zinc is sphalerite, which contains zinc sulfide (ZnS) and a small amount of cadmium sulfide (CdS).

The stages involved in the extraction of zinc from sphalerite are:


Stage 1 Sphalerite is strongly heated in air.



Stage 2 The mixture of oxides is reacted with sulfuric acid.



Stage 3 Zinc dust is added to the solution containing zinc sulfate and cadmium sulfate to remove the cadmium ions.



Stage 4 The solid cadmium is filtered off and the pure zinc sulfate solution is electrolysed.

(a) State how the reaction in stage 3 shows that zinc is more reactive than cadmium.

(1)

.....

.....


(b) (i) During the electrolysis in stage 4, zinc is deposited on the cathode.

Write an ionic half-equation for the reaction that occurs.

(1)

.....

(ii) Complete the ionic half-equation for the reaction occurring at the anode.



(1)

(iii) Explain how the pH of the solution surrounding the anode changes during the electrolysis.

(2)

.....

.....

.....

.....

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

(c) Zinc is mixed with copper to make the alloy brass.

Explain why brass is harder than pure copper.

(3)

.....

.....

.....

.....

.....

.....

.....

(Total for Question 5 = 8 marks)

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

6 The mineral nesquehonite is a form of hydrated magnesium carbonate.

The formula, $\text{MgCO}_3 \cdot x\text{H}_2\text{O}$, shows that nesquehonite contains water of crystallisation.

When a sample of nesquehonite is heated gently, the water of crystallisation is given off and anhydrous magnesium carbonate is left.

Six students are each given a sample of nesquehonite of mass 6.1 g. The students heat their samples for different times. The samples are then allowed to cool before being reweighed.


The table shows their results.

time of heating / minutes	0	1.0	2.0	3.0	4.0	4.5	6.0
mass of sample after heating / g	6.1	5.3	4.5	3.7	2.8	2.4	2.4

(a) Plot these results on the grid provided. The last two points have been plotted for you.

Draw a straight line of best fit for the points you have plotted.

(2)



- (b) Predict the mass of sample remaining after heating for 2.5 minutes.
Show on the graph how you obtain your answer. (2)

mass after 2.5 minutes = g

- (c) State why the last two masses in the table are exactly the same. (1)

- (d) A sample of nesquehonite contains 1.68 g of MgCO_3 and 1.08 g of H_2O .
Calculate the value of x in the formula $\text{MgCO}_3 \cdot x\text{H}_2\text{O}$
[M_r of $\text{MgCO}_3 = 84$; M_r of $\text{H}_2\text{O} = 18$] (3)

x =

- (e) When anhydrous magnesium carbonate is heated strongly, it decomposes.
The equation for the reaction is:



Calculate the volume of carbon dioxide formed in cm^3 , at rtp, when 4.2 g of anhydrous magnesium carbonate are completely decomposed.

[M_r of $\text{MgCO}_3 = 84$]

[Assume that the molar volume at rtp of carbon dioxide is 24 dm^3]

(3)

volume = cm^3

(Total for Question 6 = 11 marks)


DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

7 Ammonia is manufactured on a large scale and is used to make fertilisers such as ammonium nitrate (NH_4NO_3).

- (a) The first stage in the manufacture of ammonium nitrate is to react ammonia gas with oxygen gas.





The reaction is carried out at a pressure of about 10 atm and at a temperature of 800°C , in the presence of a catalyst.

If the mixture is left for long enough in a sealed container, the reaction reaches a position of dynamic equilibrium.

Graph A shows how the percentage of nitrogen monoxide (NO) in the equilibrium mixture varies with temperature at constant pressure.

Graph B shows how the percentage of nitrogen monoxide (NO) in the equilibrium mixture varies with pressure at constant temperature.



- (i) Explain why the percentage of NO at equilibrium decreases in each case.

(4)

Graph A

.....

.....

.....

Graph B

.....

.....

.....

.....

- (ii) Explain why the use of a catalyst has no effect on the position of equilibrium in a reversible reaction.

(2)


.....

.....

.....

.....

- (b) The second stage in the manufacture of ammonium nitrate is to convert the nitrogen monoxide into nitric acid. The nitric acid is then reacted with concentrated aqueous ammonia as shown in this equation.



- (i) State, in terms of proton transfer, why this reaction is classified as an acid-base reaction.

(1)

.....

.....

- (ii) Calculate the volume of 14.8 mol / dm^3 aqueous ammonia that is required to exactly neutralise 150 dm^3 of a solution of nitric acid of concentration 15.8 mol / dm^3 .

(2)

volume of aqueous ammonia = dm^3

(Total for Question 7 = 9 marks)

8 Polymers can be classified as addition polymers or condensation polymers.

(a) An addition polymer can be formed from the monomer $\text{CH}_2=\text{CHCl}$


(i) Name this monomer.

(1)

(ii) Name the addition polymer formed from this monomer.

(1)

(b) The diagram shows the repeat unit of a different addition polymer.





Draw the displayed formula of the monomer used to make this polymer.

(1)

(c) Polyesters are condensation polymers.

The structures of two monomers that are used to make a polyester are:



DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

- (i) Draw the structure of the repeat unit of the polyester formed from these two monomers.

(2)

- (ii) Identify the small molecule formed when these two monomers form the polyester.

(1)

(Total for Question 8 = 6 marks)

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

9 Grapes contain glucose that can be fermented to make ethanol.



(a) The grapes are collected and crushed to produce an aqueous solution containing glucose. Yeast is added to the solution and the mixture is left to ferment at a temperature of about 30 °C in the absence of air.

(i) State why yeast is added.

(2)

.....

.....

.....

.....

(ii) Explain why another organic compound may also be formed if fermentation is carried out in the presence of air.

(2)

.....

.....

.....

.....

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

(iii) Explain why 30°C is considered to be an optimum temperature for this reaction. (2)

.....

.....

.....

.....

(b) Ethanol can also be manufactured by reacting ethene with steam.

(i) Write the chemical equation for this reaction. (1)

.....

(ii) Name the type of reaction that occurs. (1)

.....


(iii) State **two** conditions used for this reaction in industry. (2)

1.....

2.....

(c) Grapes also contain esters.

The formula of an ester is shown.



Deduce the name of the carboxylic acid and the alcohol that can react together to make this ester. (2)

carboxylic acid.....

alcohol.....

(Total for Question 9 = 12 marks)

TOTAL FOR PAPER = 70 MARKS

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

BLANK PAGE

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

BLANK PAGE

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

BLANK PAGE