Please check the examination de	etails below	before enter	ring your can	didate information
Candidate surname			Other name	s
Pearson Edexcel International GCSE	Centre	Number		Candidate Number
Friday 11 January 2019				
Morning (Time: 2 hours)		Paper Reference 4PM0/01		
Further Pure N Paper 1	/lath	emat	tics	
Calculators may be used.				Total Marks

Instructions

- Use **black** ink or ball-point pen.
- **Fill in the boxes** at the top of this page with your name, centre number and candidate number.
- Answer **all** questions.
- Without sufficient working, correct answers may be awarded no marks.
- Answer the questions in the spaces provided
 - there may be more space than you need.

Information

- The total mark for this paper is 100.
- The marks for **each** question are shown in brackets
 - use this as a guide as to how much time to spend on each question.

Advice

- Read each question carefully before you start to answer it.
- Check your answers if you have time at the end.

Turn over ▶





Answer all TEN questions.

Write your answers in the spaces provided.

You must write down all the stages in your working.

1




Figure 1

Figure 1 shows a sector OAB of a circle, centre O.

The area of the sector is 27 cm²

The size of angle *AOB* is 1.5 radians.

Find the perimeter of the sector.

(4)

Question 1 continued

(Total for Question 1 is 4 marks)



2 The sum of the first n terms of an arithmetic series is S_n

Given that $S_n = \sum_{r=1}^n (4r+1)$

(a) show that $S_n = n(3 + 2n)$

(4)

The rth term of this arithmetic series is t_r

Given that $S_{n+3} = S_n + 3t_{15}$

(b) find the value of n.

(4)

Question 2 continued



Question 2 continued

Question 2 continued

(Total for Question 2 is 8 marks)



3 $f(x) = (2x+1)(x^2+5x-3)$

(a) Show that $f(x) = 2x^3 + 11x^2 - x - 3$

(2)

(b) Hence use algebra to solve the equation $2x^3 + 11x^2 - x - 3 = 0$

Give your roots to 3 decimal places where appropriate.

(3)

Question 3 continued

(Total for Question 3 is 5 marks)



 $\sin(A+B) = \sin A \cos B + \sin B \cos A$

$$\tan A = \frac{\sin A}{\cos A}$$

(a) Show that the equation $a \sin(x - 30)^{\circ} = b \sin(x + 30)^{\circ}$

can be written in the form $\tan x^{\circ} = \frac{a+b}{\sqrt{3}(a-b)}$

(5)




Figure 2

In triangle ABC, AC = 6 cm, BC = 14 cm, $\angle ABC = (x - 30)^{\circ}$ and $\angle BAC = (x + 30)^{\circ}$ as shown in Figure 2.

(b) Find, in degrees to 1 decimal place, the size of $\angle ACB$.

(4)

(c) Find, to 3 significant figures, the area of triangle ABC.

(2)

Question 4 continued

P 5 5 8 8 7 A 0 1 1 3 6

Question 4 continued

Question 4 continued

(Total for Question 4 is 11 marks)



5

$$f(x) = 2x^2 + 7x - 4$$

Given that f(x) can be written in the form $A(x + B)^2 + C$

(a) find the value of A, the value of B and the value of C.

(3)

- (b) Write down
 - (i) the minimum value of f(x),
 - (ii) the value of x at which this minimum occurs.

(2)

The equation f(x) = px - 6 has unequal real roots.

(c) Find the set of possible values of p.

(5)

Question 5 continued

Question 5 continued

Question 5 continued

(Total for Question 5 is 10 marks)



- 6 Given that $y = x^2 \sqrt{(2x-3)}$
 - (a) show that $\frac{dy}{dx} = \frac{x(5x 6)}{\sqrt{(2x 3)}}$

(4)

(b) find the value of $\frac{dy}{dx}$ when x = 2

(1)

The curve C has equation $y = x^2 \sqrt{(2x-3)}$

(c) Find an equation of the normal to C at the point on C where x = 2 Give your answer in the form ax + by + c = 0, where a, b and c are integers.

(5)

Question 6 continued

Question 6 continued

Question 6 continued

(Total for Question 6 is 10 marks)

7




Figure 3 shows a rectangular sheet of metal 10 cm by 16 cm. A square of side x cm is cut away from each corner of the sheet. The sheet is then folded along the dotted lines to form an open box.

The volume of the box is $V \text{ cm}^3$

(a) Show that $V = 4x^3 - 52x^2 + 160x$

(3)

(b) Using calculus, find the value of x for which V is a maximum, justifying that this value of x gives a maximum value of V.

(5)

(c) Find the maximum value of V.

(2)

Question 7 continued



Question 7 continued

24

Question 7 continued

(Total for Question 7 is 10 marks)



- 8 A curve C has equation $y = \frac{5x-3}{2x-1}$ $x \neq \frac{1}{2}$
 - (a) Write down an equation of the asymptote to C that is
 - (i) parallel to the y-axis,
 - (ii) parallel to the *x*-axis.

(2)

(b) Find the coordinates of the points of intersection of C with the coordinate axes.

(2)

(c) Using calculus show that at every point on the curve, the gradient of C is positive.

(4)


(d) Using the axes on the opposite page, sketch C, showing clearly the asymptotes and the coordinates of the points of intersection of C with the coordinate axes.

(3)

The line l is the tangent to C at the point on the curve where x = 1

(e) Find an equation of l, giving your answer in the form y = mx + c

(4)



Question 8 continued

Question 8 continued

(Total for Question 8 is 15 marks)

9 The point A has coordinates (-3, -6) and the point B has coordinates (5, -2)

The line *l* passes through the point *A* and the point *B*.

(a) Find an equation of l, giving your answer in the form y = mx + c

(3)

The point P has coordinates (k, -2). The line through A and P is perpendicular to l.

(b) Show that k = -5

(3)

The point Q has coordinates (e, f). The line through B and Q is also perpendicular to l.

Given that the length of PQ is $\sqrt{85}$ and that f > 0

(c) find the coordinates of Q.

(6)

(d) Calculate the area of quadrilateral ABQP.

(4)

Question 9 continued

Question 9 continued

Question 9 continued

(Total for Question 9 is 16 marks)



10 (a) Expand $(1-2x)^{-\frac{1}{2}}$ in ascending powers of x up to and including the term in x^3 , simplifying each term as far as possible.

(3)

(b) Write down the range of values of x for which your expansion is valid.

(1)

$$f(x) = \frac{2 - x^2}{\sqrt{(1 - 2x)}}$$

(c) Find the series expansion of f(x) in ascending powers of x up to and including the term in x^3 , simplifying each term as far as possible.


(3)

The region R is bounded by the curve with equation y = f(x), the positive x-axis, the positive y-axis and the line with equation x = 0.2

(d) Using your expansion of f(x) and algebraic integration, find an estimate for the area of R, giving your answer to 4 decimal places.

(4)

Question 10 continued



Question 10 continued

(Total for Question 10 is 11 marks)

TOTAL FOR PAPER IS 100 MARKS