



Mark Scheme (Results)

Summer 2012

GCSE Mathematics (Linear) 1MA0
Higher (Non-Calculator) Paper 1H

Edexcel and BTEC Qualifications

Edexcel and BTEC qualifications come from Pearson, the world's leading learning company. We provide a wide range of qualifications including academic, vocational, occupational and specific programmes for employers. For further information, please visit our website at www.edexcel.com.

Our website subject pages hold useful resources, support material and live feeds from our subject advisors giving you access to a portal of information. If you have any subject specific questions about this specification that require the help of a subject specialist, you may find our Ask The Expert email service helpful.

www.edexcel.com/contactus

Pearson: helping people progress, everywhere

Our aim is to help everyone progress in their lives through education. We believe in every kind of learning, for all kinds of people, wherever they are in the world. We've been involved in education for over 150 years, and by working across 70 countries, in 100 languages, we have built an international reputation for our commitment to high standards and raising achievement through innovation in education. Find out more about how we can help you and your students at: www.pearson.com/uk

Summer 2012

Publications Code UG032625

All the material in this publication is copyright

© Pearson Education Ltd 2012

NOTES ON MARKING PRINCIPLES

- 1 All candidates must receive the same treatment. Examiners must mark the first candidate in exactly the same way as they mark the last.
- 2 Mark schemes should be applied positively. Candidates must be rewarded for what they have shown they can do rather than penalised for omissions.
- 3 All the marks on the mark scheme are designed to be awarded. Examiners should always award full marks if deserved, i.e if the answer matches the mark scheme. Examiners should also be prepared to award zero marks if the candidate's response is not worthy of credit according to the mark scheme.
- 4 Where some judgement is required, mark schemes will provide the principles by which marks will be awarded and exemplification may be limited.
- 5 Crossed out work should be marked UNLESS the candidate has replaced it with an alternative response.
- 6 Mark schemes will indicate within the table where, and which strands of QWC, are being assessed. The strands are as follows:
 - i) *ensure that text is legible and that spelling, punctuation and grammar are accurate so that meaning is clear*
Comprehension and meaning is clear by using correct notation and labeling conventions.
 - ii) *select and use a form and style of writing appropriate to purpose and to complex subject matter*
Reasoning, explanation or argument is correct and appropriately structured to convey mathematical reasoning.
 - iii) *organise information clearly and coherently, using specialist vocabulary when appropriate.*
The mathematical methods and processes used are coherently and clearly organised and the appropriate mathematical vocabulary used.

7 With working

If there is a wrong answer indicated on the answer line always check the working in the body of the script (and on any diagrams), and award any marks appropriate from the mark scheme.

If working is crossed out and still legible, then it should be given any appropriate marks, as long as it has not been replaced by alternative work.

If it is clear from the working that the “correct” answer has been obtained from incorrect working, award 0 marks. Send the response to review, and discuss each of these situations with your Team Leader.

If there is no answer on the answer line then check the working for an obvious answer.

Any case of suspected misread loses A (and B) marks on that part, but can gain the M marks. Discuss each of these situations with your Team Leader.

If there is a choice of methods shown, then no marks should be awarded, unless the answer on the answer line makes clear the method that has been used.

8 Follow through marks

Follow through marks which involve a single stage calculation can be awarded without working since you can check the answer yourself, but if ambiguous do not award.

Follow through marks which involve more than one stage of calculation can only be awarded on sight of the relevant working, even if it appears obvious that there is only one way you could get the answer given.

9 Ignoring subsequent work

It is appropriate to ignore subsequent work when the additional work does not change the answer in a way that is inappropriate for the question: e.g. incorrect canceling of a fraction that would otherwise be correct

It is not appropriate to ignore subsequent work when the additional work essentially makes the answer incorrect e.g. algebra.

Transcription errors occur when candidates present a correct answer in working, and write it incorrectly on the answer line; mark the correct answer.

10 Probability

Probability answers must be given as fractions, percentages or decimals. If a candidate gives a decimal equivalent to a probability, this should be written to at least 2 decimal places (unless tenths).

Incorrect notation should lose the accuracy marks, but be awarded any implied method marks.

If a probability answer is given on the answer line using both incorrect and correct notation, award the marks.

If a probability fraction is given then cancelled incorrectly, ignore the incorrectly cancelled answer.

11 Linear equations

Full marks can be gained if the solution alone is given on the answer line, or otherwise unambiguously indicated in working (without contradiction elsewhere). Where the correct solution only is shown substituted, but not identified as the solution, the accuracy mark is lost but any method marks can be awarded.

12 Parts of questions

Unless allowed by the mark scheme, the marks allocated to one part of the question CANNOT be awarded in another.

13 Range of answers

Unless otherwise stated, when an answer is given as a range (e.g 3.5 – 4.2) then this is inclusive of the end points (e.g 3.5, 4.2) and includes all numbers within the range (e.g 4, 4.1)

Guidance on the use of codes within this mark scheme

M1 – method mark

A1 – accuracy mark

B1 – Working mark

C1 – communication mark

QWC – quality of written communication

oe – or equivalent

cao – correct answer only

ft – follow through

sc – special case

dep – dependent (on a previous mark or conclusion)

indep – independent

isw – ignore subsequent working

1MA0_1H					
Question		Working	Answer	Mark	Notes
1	(a)		Type of film Tally Frequency	2	B2 for a table with all 3 aspects: Column/row heading 'type of film' or list of at least 3 film types Column/row heading 'tally' or tally marks (or key) Column/row heading 'frequency' or totals oe (B1 for a table with 2 of the 3 aspects)
	(b)			1	B1 for acceptable reason eg. all same age, sample too small, biased, same school

1MA0_1H					
Question	Working	Answer	Mark	Notes	
2	(a)	$360 \div 60 = 6$ $300 \div 60 = 5$ $6 \times 5 =$	Yes and 30	3	<p>M1 for dividing side of patio by side of paving slab eg. $360 \div 60$ or $300 \div 60$ or $3.6 \div 0.6$ or $3 \div 0.6$ or 6 and 5 seen (may be on a diagram) or 6 divisions seen on length of diagram or 5 divisions seen on width of diagram M1 for correct method to find number of paving slabs eg. $(360 \div 60) \times (300 \div 60)$ oe or 6×5 or 30 squares seen on diagram (units may not be consistent) A1 for Yes and 30 (or 2 extra) with correct calculations</p> <p>OR M1 for correct method to find area of patio or paving slab eg 360×300 or 108000 seen or 60×60 or 3600 seen or 3.6×3 or 10.8 seen or 0.6×0.6 or 0.36 seen M1 for dividing area of patio by area of a paving slab eg. $(3.6 \times 3) \div (0.6 \times 0.6)$ oe (units may not be consistent) A1 for Yes and 30 (or 2 extra) with correct calculations</p> <p>OR M1 for method to find area of patio or area of 32 slabs eg. $60 \times 60 \times 32$ or 360×300 M1 for method to find both area of patio and area of 32 slabs eg. $60 \times 60 \times 32$ and 360×300 (units may not be consistent) A1 for Yes and 115200 and 108000 OR Yes and 11.52 and 10.8</p> <p>NB : Throughout the question, candidates could be working in metres or centimetres</p>

1MA0_1H																
Question	Working	Answer	Mark	Notes												
(b)	$\begin{array}{r} 1726 \\ 25890 \\ \hline 27616 \end{array}$ <table border="1"> <tbody> <tr> <td></td> <td>800</td> <td>60</td> <td>3</td> </tr> <tr> <td>30</td> <td>24000</td> <td>1800</td> <td>90</td> </tr> <tr> <td>2</td> <td>1600</td> <td>120</td> <td>6</td> </tr> </tbody> </table> $24000+1800+90+1600+120+6 = 27616$		800	60	3	30	24000	1800	90	2	1600	120	6	276.16	3	<p>M1 for complete correct method with relative place value correct. Condone 1 multiplication error, addition not necessary.</p> <p>OR</p> <p>M1 for a complete grid. Condone 1 multiplication error, addition not necessary.</p> <p>OR</p> <p>M1 for sight of a complete partitioning method, condone 1 multiplication error. Final addition not necessary.</p> <p>A1 for digits 27616</p> <p>A1 ft (dep on M1) for correct placement of decimal point after addition (of appropriate values)</p> <p>(SC: B1 for attempting to add 32 lots of 8.63)</p>
	800	60	3													
30	24000	1800	90													
2	1600	120	6													

1MA0_1H		Working	Answer	Mark	Notes																					
Question																										
3	(a)		10	1	B1 cao																					
	(b)		<p>Ed is cheaper up to 20 miles, Bill is cheaper for more than 20 miles</p> <table border="1"> <thead> <tr> <th>Miles</th> <th>0</th> <th>10</th> <th>20</th> <th>30</th> <th>40</th> <th>50</th> </tr> </thead> <tbody> <tr> <td>Ed</td> <td>0</td> <td>15</td> <td>30</td> <td>45</td> <td>60</td> <td>75</td> </tr> <tr> <td>Bill</td> <td>10</td> <td>20</td> <td>30</td> <td>40</td> <td>50</td> <td>60</td> </tr> </tbody> </table>	Miles	0	10	20	30	40	50	Ed	0	15	30	45	60	75	Bill	10	20	30	40	50	60	3	3
Miles	0	10	20	30	40	50																				
Ed	0	15	30	45	60	75																				
Bill	10	20	30	40	50	60																				

1MA0_1H					
Question	Working	Answer	Mark	Notes	
4	$\begin{array}{r l} 2 & 9 \\ 3 & 1\ 3\ 5\ 6\ 9 \\ 4 & 2\ 3\ 3\ 4\ 6\ 8\ 9 \\ 5 & 2\ 4\ 5 \end{array}$ <p>OR</p> $\begin{array}{r l} 20 & 9 \\ 30 & 1\ 3\ 5\ 6\ 9 \\ 40 & 2\ 3\ 3\ 4\ 6\ 8\ 9 \\ 50 & 2\ 4\ 5 \end{array}$	$\begin{array}{r l} 2 & 9 \\ 3 & 1\ 3\ 5\ 6\ 9 \\ 4 & 2\ 3\ 3\ 4\ 6\ 8\ 9 \\ 5 & 2\ 4\ 5 \end{array}$ <p>Key: $2 \mid 9 = 29$</p>	3	<p>B3 for fully correct diagram with appropriate key (B2 for ordered leaves, with at most two errors or omissions and a key OR correct unordered leaves and a key OR correct ordered leaves)</p> <p>(B1 for unordered or ordered leaves, with at most two errors or omissions OR key)</p> <p>NB : Order of stem may be reversed; condone commas between leaves</p>	
5	$c = \frac{30 \times 40}{150}$	8	2	<p>M1 for $\frac{30 \times 40}{150}$ or 1200 seen A1 cao</p>	
6	(a)	30	2	<p>M1 for $25 \div 10$ or 2.5 seen or $10 \div 25$ or 0.4 seen or $12 + 12 + 6$ oe or a complete method eg. $25 \times 12 \div 10$ oe A1 cao</p>	
	(b)	$1000 \div 200 \times 12$	60	2	<p>M1 for $500 \div 50$ or $1000 \div 200$ or $500 \div 10$ OR correct scale factor clearly linked with one ingredient eg. 10 with sugar or 5 with butter or flour or 50 with milk OR answer of 120 or 600 A1 cao</p>

1MA0_1H				
Question	Working	Answer	Mark	Notes
7	<p>Acton after 24, 48, 72, 96, 120 Barton after 20, 40, 60, 80, 100, 120 LCM of 20 and 24 is 120 9:00 am + 120 minutes</p> <p>OR Acton after 24, 48, 1h 12 m, 1h 36m, 2h Barton after 20, 40, 1 h, 1h 20m, 1h 40m, 2h LCM is 2 hours 9:00 am + 2 hours</p> <p>OR Times from 9:00 am when each bus leaves the bus station Acton at 9:24, 9:48, 10:12, 10:36, 11:00 Barton at 9:20, 9:40, 10:00, 10:20, 10:40, 11:00</p> <p>OR $20 = 2 \times 2 \times 5$ $24 = 2 \times 2 \times 2 \times 3$ $2 \times 2 \times 2 \times 3 \times 5 = 120$</p>	11:00 am	3	<p>M1 for listing multiples of 20 and 24 with at least 3 numbers in each list ; multiples could be given in minutes or in hours and minutes (condone one addition error in total in first 3 numbers in lists) A1 identify 120 (mins) or 2 (hours) as LCM A1 for 11:00 (am) or 11(am) or 11 o'clock</p> <p>OR M1 for listing times after 9am when each bus leaves the bus station, with at least 3 times in each list (condone one addition error in total in first 3 times after 9am in lists) A1 for correct times in each list up to and including 11:00 A1 for 11:00 (am) or 11(am) or 11 o'clock</p> <p>OR M1 for correct method to write 20 and 24 in terms of their prime factors 2, 2, 5 and 2, 2, 2, 3 (condone one error) A1 identify 120 as LCM A1 for 11:00 (am) or 11(am) or 11 o'clock</p>

1MA0_1H				
Question	Working	Answer	Mark	Notes
8	(a)	$6y - 15$	1	B1 cao
	(b)	$4x(2x + y)$	2	B2 cao (B1 for $x(8x + 4y)$ or $2x(4x + 2y)$ or $4(2x^2 + xy)$ or $4x(ax + by)$ where a, b are positive integers or $ax(2x + y)$ where a is a positive integer or $4x(2x - y)$)
	(c)	$\frac{10t}{g}$	2	M1 for clear intention to multiply both sides of the equation by 10 (eg. $\times 10$ seen on both sides of equation) or clear intention to divide both sides of the equation by g (eg. $\div g$ seen on both sides of equation) or $10t = gh$ or $\frac{t}{g} = \frac{h}{10}$ or fully correct reverse flow diagram eg. $\leftarrow \times 10 \leftarrow \div g \leftarrow$ A1 for $\frac{10t}{g}$ oe

1MA0_1H					
Question		Working	Answer	Mark	Notes
9			Rotation 180° Centre (3, 3) or Enlargement Scale factor -1 Centre (3, 3)	3	B1 for rotation B1 for 180° B1 for (3, 3) OR B1 for enlargement B1 for scale factor -1 B1 for (3, 3) B0 for a combination of transformations


1MA0_1H				
Question	Working	Answer	Mark	Notes
10	$2.25 \times 60 \div 100 = 1.35$ $1.35 + 0.80 = 2.15$ $1.5 \times 60 \div 100 = 0.90$ $0.90 + 1.90 = 2.80$	Railtickets with correct calculations	4	<p>NB. All work may be done in pence throughout</p> <p>M1 for correct method to find credit card charge for one company eg. $0.0225 \times 60 (=1.35)$ oe or $0.015 \times 60 (=0.9)$ oe M1 (dep) for correct method to find total additional charge or total price for one company eg. $0.0225 \times 60 + 0.80$ or $0.015 \times 60 + 1.90$ or 2.15 or $2.8(0)$ or 62.15 or $62.8(0)$ A1 for 2.15 and $2.8(0)$ or 62.15 and $62.8(0)$ C1 (dep on M1) for a statement deducing the cheapest company, but figures used for the comparison must also be stated somewhere, and a clear association with the name of each company</p> <p>OR</p> <p>M1 for correct method to find percentage of (60+booking fee) eg. $0.0225 \times 60.8 (=1.368)$ oe or $0.015 \times 61.9 (=0.9285)$ M1 (dep) for correct method to find total cost or total additional cost eg. '1.368' + $60.8 (=62.168)$ or '1.368' + $0.8 (=2.168)$ or '0.9285' + $61.9 (=62.8285)$ or '0.9285' + $1.9 (=2.8285)$ A1 for 62.168 or 62.17 AND 62.8285 or 62.83 OR 2.168 or 2.17 AND 2.8285 or 2.83 C1 (dep on M1) for a statement deducing the cheapest company, but figures used for the comparison must also be stated somewhere, and a clear association with the name of each company</p> <p>OR</p>
	OR			OR

1MA0_1H				
Question	Working	Answer	Mark	Notes
	$2.25 - 1.5 = 0.75$ $0.075 \times 60 \div 100 = 0.45$ $0.80 + 0.45 = 1.25$ $1.25 < 1.90$			<p>M1 for correct method to find difference in cost of credit card charge eg. $(2.25 - 1.5) \times 60 \div 100$ or 0.45 seen M1 (dep) for using difference with booking fee or finding difference between booking fees eg. $0.80 + "0.45" (=1.25)$ or $1.90 - "0.45" (=1.45)$ or $1.90 - 0.8 (=1.1(0))$ A1 1.25 and 1.9(0) or 0.45 and 1.1(0) C1 (dep on M1) for a statement deducing the cheapest company, but figures used for the comparison must also be stated somewhere, and a clear association with the name of each company</p> <p>QWC: Decision and justification should be clear with working clearly presented and attributable</p>

1MA0_1H				
Question	Working	Answer	Mark	Notes
11	$3x-15 = 2x+24$ $x = 39$ OR $2x+3x-15 +2x+ 2x+24 = 360$ $9x + 9 = 360$ $9x = 351$ $x = 39$ OR $2x + 2x+24 = 180$ $4x + 24 = 180$ $4x = 156$ $x = 39$ OR $2x + 3x-15 = 180$ $5x - 15 = 180$ $5x = 195$ $x = 39$	39	3	<p>M1 for forming an appropriate equation eg. $3x - 15 = 2x + 24$</p> <p>OR $2x + 3x - 15 + 2x + 2x + 24 = 360$</p> <p>OR $2x + 2x + 24 = 180$</p> <p>OR $2x + 3x - 15 = 180$</p> <p>OR $2x + 3x - 15 = 2x + 2x + 24$</p> <p>M1 (dep) for correct operation(s) to isolate x and non-x terms in an equation to get to $ax = b$ A1 cao</p> <p>OR M2 for $\frac{351}{9}$ oe or $\frac{195}{5}$ oe or $\frac{156}{4}$ oe A1 cao</p>

1MA0_1H				
Question	Working	Answer	Mark	Notes
12	$6 \times 10 \times 8 = 480$ $480 \div (6 \times 20) =$	4	3	<p>M1 for $6 \times 10 \times 8$ or 480 seen M1 (dep) for '480' $\div (6 \times 20)$ oe A1 cao</p> <p>OR</p> <p>M1 for $20 \div 10 (=2)$ or $10 \div 20 (= \frac{1}{2})$ or $\frac{8}{20}$ oe or $\frac{20}{8}$ oe</p> <p>M1 (dep) for $8 \div '2'$ or $8 \times \frac{1}{2}$ or $\frac{8}{20} \times 10$ oe or</p> <p>$10 \div \frac{20}{8}$</p> <p>A1 cao</p> <p>SC : B2 for answer of 16 coming from $\frac{20 \times 8 \times 6}{10 \times 6}$ oe</p>

1MA0_1H				
Question	Working	Answer	Mark	Notes
13	$180 - (360 \div 6) = 120$ $180 - (360 \div 8) = 135$ $360 - 120 - 135 =$ OR $360 \div 6 = 60$ $360 \div 8 = 45$ $60 + 45 =$	105	4	<p>NB. Do remember to look at the diagram when marking this question. Looking at the complete method should confirm if interior or exterior angles are being calculated</p> <p>M1 for a correct method to work out the interior angle of a regular hexagon eg. $180 - (360 \div 6)$ oe or $(6 - 2) \times 180 \div 6$ oe or 120 as interior angle of the hexagon M1 for a correct method to work out the interior angle of a regular octagon $180 - (360 \div 8)$ oe or $(8 - 2) \times 180 \div 8$ oe or 135 as interior angle of the octagon M1 (dep on at least M1) for a complete method eg. $360 - "120" - "135"$ A1 cao</p> <p>OR M1 for a correct method to work out an exterior angle of a regular hexagon eg. $360 \div 6$ or 60 as exterior angle of the hexagon M1 for a correct method to work out an exterior angle of a regular hexagon $360 \div 8$ or 45 as exterior angle of the octagon M1 (dep on at least M1) for a complete method eg. $"60" + "45"$ A1 cao</p> <p>SC : B1 for answer of 255</p>

1MA0_1H				
Question	Working	Answer	Mark	Notes
14	(a)	35	1	B1 for 34 – 36
	(b)	110	1	B1 for 108 – 112
	(c)	Position of <i>B</i> marked	2	B1 for a point marked on a bearing of $40^\circ (\pm 2^\circ)$ from <i>H</i> or for a line on a bearing of $40^\circ (\pm 2^\circ)$ (use straight line guidelines on overlay) B1 for a point 4 cm (± 0.2 cm) from <i>H</i> or for a line of length 4 cm (± 0.2 cm) from <i>H</i> (use circular guidelines on overlay) NB. No label needed for point
15	(a)	170	1	B1 accept answers in range 170 - 170.5 inclusive
	(b)		3	B3 for box plot with all 3 aspects correct (overlay) aspect 1 : ends of whiskers at 153 and 186 aspect 2 : ends of box at 165 and 175 aspect 3 : median marked at 170 or ft (a) provided $165 < (a) < 175$ (B2 for box plot with two aspects correct) (B1 for one aspect or correct quartiles and median identified) SC : B2 for all 5 values (153, 165, '170', 175, 186) plotted
	(c)	Two correct comparisons	2	B1 ft from (b) for a correct comparison of range or inter-quartile range eg. the range / iqr is smaller for group B than group A B1 ft from (b) for a correct comparison of median or upper quartile or lower quartile or minimum or maximum eg. the median in group A is greater than the median in group B

1MA0_1H				
Question	Working	Answer	Mark	Notes
16	(a)		1	B1 for m^{-10} or $\frac{1}{m^{10}}$
	(b)		2	M1 for $(x \pm 5)(x \pm 2)$ or $x(x - 2) + 5(x - 2)$ or $x(x + 5) - 2(x + 5)$ A1
17	(a)		1	B1 cao
	(b)		1	B1 cao
	(c)		2	M1 for $27 \times 10^{7+6}$ or 27×10^{13} oe or an answer of 2.7×10^n where n is an integer or an answer of $a \times 10^{14}$ where $1 \leq a < 10$ A1 cao

1MA0_1H				
Question	Working	Answer	Mark	Notes
18	$\frac{1}{2} \times 4 \times 3 = 6$ $\left(\frac{1}{2}\right)^2 \times 6 =$	1.5	3	<p>M1 for $\frac{1}{2} \times 4 \times 3$ oe</p> <p>M1 for $\left(\frac{1}{2}\right)^2 \times "6"$</p> <p>A1 cao</p> <p>OR</p> <p>M2 for $\frac{1}{2} \times 2 \times 1.5$ oe</p> <p>(M1 for triangle with all lengths $\frac{1}{2}$ corresponding lengths of triangle <i>ABC</i> seen in any position or vertices seen at (1, 1) (3,1) and (2.5, 2.5) or stated)</p> <p>A1 cao</p>
19	(a)	0.6 0.7, 0.3, 0.7	2	<p>B1 for 0.6 in correct position on tree diagram</p> <p>B1 for 0.7, 0.3, 0.7 in correct positions on tree diagram</p>
	(b)	$0.4 \times 0.3 =$	2	<p>M1 for 0.4×0.3 oe or a complete alternative method ft from tree diagram</p> <p>A1 for 0.12 oe</p>

1MA0_1H				
Question	Working	Answer	Mark	Notes
20	$15x + 6y = 33$ $8x - 6y = 36$ $23x = 69$ $5 \times 3 + 2y = 11$ OR $x = \frac{11-2y}{5}$ $4 \times \left(\frac{11-2y}{5} \right) - 3y = 18$ $44 - 8y - 15y = 90$ $-46 = 23y$ $y = -2$	$x = 3$ $y = -2$	4	<p>M1 for coefficients of x or y the same followed by correct operation (condone one arithmetic error) A1 cao for first solution M1 (dep on M1) for correct substitution of found value into one of the equations or appropriate method after starting again (condone one arithmetic error) A1 cao for second solution</p> <p>OR M1 for full method to rearrange and substitute to eliminate x or y, (condone one arithmetical error) A1 cao for first solution M1 (dep on M1) for correct substitution of found value into one of the equations or appropriate method after starting again (condone one arithmetic error) A1 cao for second solution</p> <p>Trial and improvement 0 marks unless both x and y correct values found</p>

1MA0_1H				
Question	Working	Answer	Mark	Notes
21*	<p>$ABO = ADO = 90^\circ$ (Angle between tangent and radius is 90°) $DOB = 360 - 90 - 90 - 50$ (Angles in a quadrilateral add up to 360°) $BCD = 130 \div 2$ (Angle at centre is twice angle at circumference)</p> <p>OR $ABD = (180 - 50) \div 2$ (Base angles of an isosceles triangle) $BCD = 65$ (Alternate segment theorem)</p>	65°	4	<p>B1 for $ABO = 90$ or $ADO = 90$ (may be on diagram) B1 for $BCD = 65$ (may be on diagram)</p> <p>C2 for $BCD = 65^\circ$ stated or $DCB = 65^\circ$ stated or angle C = 65° stated with all reasons: angle between <u>tangent</u> and <u>radius</u> is 90°; <u>angles in a quadrilateral</u> sum to 360°; <u>angle at centre</u> is <u>twice</u> angle at <u>circumference</u></p> <p>(accept angle at circumference is half (or $\frac{1}{2}$) the angle at the centre)</p> <p>(C1 for one correct and appropriate circle theorem reason) QWC: Working clearly laid out and reasons given using correct language</p> <p>OR B1 for $ABD = 65$ or $ADB = 65$ (may be on diagram) B1 for $BCD = 65$ (may be on diagram)</p> <p>C2 for $BCD = 65^\circ$ stated or $DCB = 65^\circ$ stated or angle C = 65° stated with all reasons: base <u>angles</u> of an <u>isosceles triangle</u> are <u>equal</u>; <u>angles in a triangle</u> sum to 180°; <u>tangents from an external point</u> are <u>equal</u>; <u>alternate segment</u> theorem</p> <p>(C1 for one correct and appropriate circle theorem reason) QWC: Working clearly laid out and reasons given using correct language</p>

1MA0_1H																		
Question	Working				Answer	Mark	Notes											
22	(a)	<table border="1"> <tr> <td>F</td> <td>15</td> <td>25</td> <td>36</td> <td>24</td> </tr> <tr> <td>Fd</td> <td>3</td> <td>5</td> <td>3.6</td> <td>1.2</td> </tr> </table>				F	15	25	36	24	Fd	3	5	3.6	1.2	Correct histogram	3	<p>B3 for fully correct histogram (overlay) (B2 for 3 correct blocks) (B1 for 2 correct blocks of different widths)</p> <p>SC : B1 for correct key, eg. $1 \text{ cm}^2 = 5 \text{ (cars)}$ or correct values for (freq \div class interval) for at least 3 frequencies (3, 5, 3.6, 1.2)</p> <p>NB: The overlay shows one possible histogram, there are other correct solutions.</p>
	F	15	25	36	24													
Fd	3	5	3.6	1.2														
(b)	$\frac{3}{4} \times 24$				18	2	<p>M1 for $\frac{3}{4} \times 24 (=18)$ oe or $\frac{1}{4} \times 24 (=6)$ oe A1 cao</p> <p>OR</p> <p>M1 ft histogram for $15 \times "1.2"$ or $5 \times "1.2"$ A1 ft</p>											

1MA0_1H				
Question	Working	Answer	Mark	Notes
23	(a)	$\frac{(x+4)(x-1)}{(2x-3)(x-1)}$	$\frac{x+4}{2x-3}$	3 M1 for $(x+4)(x-1)$ M1 for $(2x-3)(x-1)$ A1 cao
	(b)	$\frac{4(x-2)}{(x+2)(x-2)} + \frac{3(x+2)}{(x+2)(x-2)}$	$\frac{7x-2}{(x+2)(x-2)}$	3 M1 for denominator $(x+2)(x-2)$ oe or x^2-4 M1 for $\frac{4(x-2)}{(x+2)(x-2)}$ oe or $\frac{3(x+2)}{(x+2)(x-2)}$ oe (NB. The denominator must be $(x+2)(x-2)$ or x^2-4 or another suitable common denominator) A1 for $\frac{7x-2}{(x+2)(x-2)}$ or $\frac{7x-2}{x^2-4}$ SC: If no marks awarded then award B1 for $\frac{4(x-2)}{x^2-2} + \frac{3(x+2)}{x^2-2}$ oe

1MA0_1H				
Question	Working	Answer	Mark	Notes
24	eg. $x = 0.28181\dots$ $100x = 28.181\dots$ $99x = 27.9$	$\frac{31}{110}$	3	M1 for 0.28181(...) or $0.2 + 0.08181\dots$ or evidence of correct recurring decimal eg. 281.81(...) M1 for two correct recurring decimals that, when subtracted, would result in a terminating decimal, and attempting the subtraction eg. $100x = 28.1818\dots$, $x = 0.28181\dots$ and subtracting eg. $1000x = 281.8181\dots$, $10x = 2.8181\dots$ and subtracting OR $\frac{27.9}{99}$ or $\frac{279}{990}$ oe A1 cao
25	$\text{Vol cylinder} = \pi \times (2x)^2 \times 9x$ $= 36\pi x^3$ $36\pi x^3 = \frac{4}{3}\pi r^3$ $r^3 = 27x^3$	$3x$	3	M1 for sub. into $\pi r^2 h$ eg. $\pi \times (2x)^2 \times 9x$ oe M1 for $\pi \times (2x)^2 \times 9x = \frac{4}{3}\pi r^3$ oe A1 oe eg. $\sqrt[3]{\frac{36x^3}{4}} \frac{4}{3}$ NB : For both method marks condone missing brackets around the $2x$

1MA0_1H					
Question		Working	Answer	Mark	Notes
26	(a)		Parabola through (4, -1), (2, 3), (6, 3) (3, 0) (5, 0)	2	B2 for a parabola with min (4, -1), through (2, 3), (6, 3), (3, 0), (5, 0) (B1 for a parabola with min (4, -1) or a parabola through (2, 3) and (6, 3) or a parabola through (3, 0) and (5, 0) or a translation of the given parabola along the x -axis by any value other than +3 with the points (-1, 3) (0, 0) (1, -1) (2, 0) (3, 3) all translated by the same amount)
	(b)		Parabola through (1, -2), (0, 0), (2, 0)	2	B2 parabola with min (1, -2), through (0, 0) and (2, 0) (B1 parabola with min (1, -2) or parabola through (0, 0), (2, 0) (-1, 6) and (3, 6))

Further copies of this publication are available from
Edexcel Publications, Adamsway, Mansfield, Notts, NG18 4FN

Telephone 01623 467467

Fax 01623 450481

Email publication.orders@edexcel.com

Order Code UG032625 Summer 2012

For more information on Edexcel qualifications, please visit our website
www.edexcel.com

Pearson Education Limited. Registered company number 872828
with its registered office at Edinburgh Gate, Harlow, Essex CM20 2JE

Ofqual




Llywodraeth Cynulliad Cymru
Welsh Assembly Government

