Chemistry 2 - Common questions

Que Nur	stion nber						
FT	HT	Sub-section	Mark	Answer	Accept	Neutral answer	Do not accept
8	1	(a)	1	solubility of sodium carbonate increases (until 40-43°C) and then decreases			
		(b)	3	plotting 7 correct points(2)plotting 6 correct points(1)suitable line – must be curve(1)			
		(c)	1	sodium carbonate sodium bromate sodium chloride – correct order			
		(d)	3	recognise that sodium chloride is soluble and silver chloride is notadd (enough/excess) water (to remove/dissolve all the sodium chloride)(1)filter(1)			

Question Number									
FT	ΗT	Su	b-sect	ion	Mark	Answer	Accept	Neutral answer	Do not accept
9	2	(a)	(i)		2	436 + 242 (1)			·
						= 678 (1) - correct answer only (cao) (2)			
	<u> </u>		(ii)		2	$2 \times 431 (1)$ = 862 (1) - cao (2)			
		(b)			1	exothermic since energy given out (as bonds made) > energy needed (to break the bonds) energy given by reaction is negative / -184 credit 'endothermic' with correct reason if calculation error followed through (ft)			

Su	ub-section Mark Answer				Neutral	Do not accept
					answer	
(a)	(i)	1	A – requires the most soap – both needed			
	(ii)	2	D(1)some hardness removed by boiling but not all / temporary hardness removed by boiling but permanent hardness remained(1)Alternative answer accepted for all candidates due to very common mis-interpretation of question on Welsh-medium papersA contains permanent hardness and C contains temporary hardness(1)A loses none of its hardness through boiling and C loses all of its hardness (through boiling)			
(b)		2	same trend / A still the hardest / B still the softest / D still contains both permanent and temporary hard water (1) different amount of water used / different concentration of soap solution / shaken for a different amount of time / different amount of lather formed (1)			
	· Su (a)	· Sub-secti (a) (i) (ii) (iii) (b) (b)	Sub-section Mark (a) (i) 1 (ii) 2 (iii) 2 (b) 2	Sub-section Mark Answer (a) (i) 1 A – requires the most soap – both needed (a) (ii) 2 D (1) (iii) 2 D (1) some hardness removed by boiling but not all / temporary hardness removed by boiling but permanent hardness remained (1) Alternative answer accepted for all candidates due to very common mis-interpretation of question on Welsh-medium papers A contains permanent hardness and C contains temporary hardness A loses none of its hardness through boiling and C loses all of its hardness (through boiling) (1) (b) 2 same trend / A still the hardest / B still the softest / D still contains both permanent and temporary hard water (1) different amount of water used / different concentration of soap solution / shaken for a different amount of time / different amount of lather formed	Sub-section Mark Answer Accept (a) (i) 1 A – requires the most soap – both needed Accept (a) (ii) 2 D (1) (iii) 2 D (1) some hardness removed by boiling but not all / temporary hardness removed by boiling but permanent hardness remained (1) Alternative answer accepted for all candidates due to very common mis-interpretation of question on Welsh-medium papers A contains permanent hardness and C contains temporary hardness A loses none of its hardness through boiling and C loses all of its hardness (through boiling) (1) (b) 2 same trend / A still the hardest / B still the softest / D still contains both permanent and temporary hard water (1) different amount of water used / different concentration of soap solution / shaken for a different amount of time / different amount of lather formed	Sub-section Mark Answer Accept Neutral answer (a) (i) 1 A – requires the most soap – both needed

Que	stion		
FT	HT	Mark	Guidance
11	4	6	 Indicative content: This method of separation is called fractional distillation. Crude oil is a mixture of hydrocarbons. The crude oil is heated and vaporised before entering the column. Smaller/lower boiling hydrocarbons will rise in the column and condense higher up the column. Hydrocarbons with similar boiling points condense at the same level in the column. (Boiling point depends on the size of the molecule – larger molecules have higher boiling points.) 5-6 marks: The candidate constructs an articulate, integrated account correctly linking relevant points, such as those in the indicative content, which shows sequential reasoning. The answer fully addresses the question with no irrelevant inclusions or significant omissions. The candidate uses appropriate scientific terminology and accurate spelling, punctuation and grammar.
			 3-4 marks: The candidate constructs an account correctly linking some relevant points, such as those in the indicative content, showing some reasoning. The answer addresses the question with some omissions. The candidate uses mainly appropriate scientific terminology and some accurate spelling, punctuation and grammar. 1-2 marks: The candidate makes some relevant points, such as those in the indicative content, showing limited reasoning. The answer addresses the question with significant omissions. The candidate uses limited scientific terminology and inaccuracies in spelling, punctuation and grammar. 0 marks: The candidate does not make any attempt or give a relevant answer worthy of credit.

Chemistry 2 - Higher Tier only questions

Que	stion								
Nun	1ber				AA 1.				
F 1	HI	Su	D-Secti	ion /	Mark	Answer	Accept	Neutral answer	Do not accept
	5	(a)			I	2,8,1			
		(b)			2	$\begin{array}{c} \begin{array}{c} \begin{array}{c} \begin{array}{c} \end{array} \\ \end{array} \\ \end{array} \\ \begin{array}{c} \end{array} \\ \end{array} \\ \end{array} \\ \begin{array}{c} \end{array} \\ \end{array} \\ \begin{array}{c} \end{array} \\ \end{array} \\ \end{array} \\ \end{array} \\ \begin{array}{c} \end{array} \\ \end{array} \\ \end{array} \\ \end{array} \\ \begin{array}{c} \end{array} \\ \end{array} $			
		(c)	(i)		1	floats moves fizzes / bubbles goes into a round shape / melts – any two		vigorous reaction dissolves	
			(ii)		1	sodium hydroxide and hydrogen – both needed	NaOH + H ₂	Н	
		(d)			1	potassium burns / lilac flame		potassium moves faster	yellow / orange / red / green flame
		(e)			2	atoms get bigger / greater distance between the (positive) nucleus and the (outer) electron(1)outer electron more weakly held(1)			

_

Que: Num	stion Iber								
FT	HT	Su	b-sect	ion	Mark	Answer	Accept	Neutral answer	Do not accept
	6	(a) (i)			1	polytetrafluoroethene / PTFE		Teflon	
			(ii)	Ι	1	$ \begin{array}{c c} & CH_3 & H \\ & I \\ & C \\ & C \\ & C \\ & H \\ & H \\ & n \\ \end{array} $			
				II	1	addition			additional
		(b)			2	both have long or large molecules / long chains of carbon atoms / polymer chains(1)			
						(only) thermosets have crosslinking / strong bonds between chains (1)			reference to layers

Que Nur	stion nber								
FT	HT	Su	b-sect	ion	Mark	Answer	Accept	Neutral answer	Do not accept
	7	(a)			1	lithium / Li ⁺	Li Ca Sr		
	•	(b)	(i)		1	cream precipitate	off white ppt		pale yellow ppt
			(ii)		2	$\begin{array}{c} Ag^{+} + Br^{-} \qquad (1) \\ AgBr \qquad (1) \end{array}$			
		(c)	(i)		1	chlorine / fluorine	Cl ₂ / F ₂	Cl / F	
			(ii)		1	displacement	redox		

Question Number									
FT	HT	Su	b-sectio	on Mark	Answer		Accept	Neutral answer	Do not accept
	8	(a)		1	any one from: some copper had not reacted reaction had not been completed heated to constant mass				
		(b)		3	mass of copper = $37.7 - 25.0 = 12.7$ mass of oxygen = $40.9 - 37.7 = 3.2$ n(Cu) = $12.7/63.5$ n(O) = $3.2/16$ ratio $0.2/0.2 : 0.2/0.2$ ie 1:1 formula CuO	 (1) (1) (1) 			

Que Nun	stion nber								
FT	HT	Su	b-sect	ion N	Nark	Answer	Accept	Neutral answer	Do not accept
	9	(a)			3	two discrete diagrams needed: - diagram 1 showing transfer of electrons - diagram 2 showing ions			
						diagram 1two lithium atoms losing 1 electron eachone sulfur atom gaining 2 electrons(1)			
						diagram 2 $2 \times \text{Li}^+$ and S ²⁻ formed (1)			
		(b)			2	magnesium and/or oxide ions have a greater charge than sodium and/or chloride ions (1)this gives a greater attraction between the ions / stronger ionic bonds / more energy is needed to break bonds (1)			
		(c)			2	shared electron pair between H atoms and adjacent O atoms and between the two O atoms(1)8 electrons in outer shell of both O atoms(1)			

Ques	tion		
Numb	oer		
FT	ΗT	Mark	Guidance
	10	6	 Indicative content: For a chemical reaction to take place the reactant particles must collide. Increasing the concentration increases the number of particles in the same volume which gives a greater chance of the particles colliding, giving an increase in the rate of the reaction. As the temperature increases the reactant particles are moving faster, increasing the chance of a collision. At higher temperature the particles also have higher energy which increases the possibility of having sufficient energy during collision to overcome the activation energy and become a 'successful collision'. Therefore increasing the temperature also increases the speed of a reaction. 5-6 marks: The candidate constructs an articulate, integrated account correctly linking relevant points, such as those in the indicative content, which shows sequential reasoning. The answer fully addresses the question with no irrelevant inclusions or significant omissions. The candidate uses appropriate scientific terminology and accurate spelling, punctuation and grammar. 3-4 marks: The candidate constructs an account correctly linking some relevant points, such as those in the indicative content, showing some accurate spelling, punctuation and grammar. 1-2 marks: The candidate makes some relevant points, such as those in the indicative content, showing limited reasoning. The candidate uses appropriate scientific terminology and inaccuracies in spelling, punctuation and grammar. 0 marks: The candidate does not make any attempt or give a relevant answer worthy of credit.