



Mark Scheme (Results)

November 2020

Pearson Edexcel International GCSE Mathematics A (4MA1) Paper 2F

Edexcel and BTEC Qualifications

Edexcel and BTEC qualifications are awarded by Pearson, the UK's largest awarding body. We provide a wide range of qualifications including academic, vocational, occupational and specific programmes for employers. For further information visit our qualifications websites at <u>www.edexcel.com</u> or <u>www.btec.co.uk</u>. Alternatively, you can get in touch with us using the details on our contact us page at <u>www.edexcel.com/contactus</u>.

Pearson: helping people progress, everywhere

Pearson aspires to be the world's leading learning company. Our aim is to help everyone progress in their lives through education. We believe in every kind of learning, for all kinds of people, wherever they are in the world. We've been involved in education for over 150 years, and by working across 70 countries, in 100 languages, we have built an international reputation for our commitment to high standards and raising achievement through innovation in education. Find out more about how we can help you and your students at: <u>www.pearson.com/uk</u>

November 2020 Publications Code 4MA1_2F_2011_MS All the material in this publication is copyright © Pearson Education Ltd 2021 **General Marking Guidance**

- All candidates must receive the same treatment. Examiners must mark the first candidate in exactly the same way as they mark the last.
- Mark schemes should be applied positively. Candidates must be rewarded for what they have shown they can do rather than penalised for omissions.
- Examiners should mark according to the mark scheme not according to their perception of where the grade boundaries may lie.
- There is no ceiling on achievement. All marks on the mark scheme should be used appropriately.
- All the marks on the mark scheme are designed to be awarded. Examiners should always award full marks if deserved, i.e. if the answer matches the mark scheme.

Examiners should also be prepared to award zero marks if the candidate's response is not worthy of credit according to the mark scheme.

- Where some judgement is required, mark schemes will provide the principles by which marks will be awarded and exemplification may be limited.
- When examiners are in doubt regarding the application of the mark scheme to a candidate's response, the team leader must be consulted.
- Crossed out work should be marked UNLESS the candidate has replaced it with an alternative response.

• Types of mark

- M marks: method marks
- A marks: accuracy marks
- B marks: unconditional accuracy marks (independent of M marks)

• Abbreviations

- o cao correct answer only
- o ft follow through
- o isw ignore subsequent working
- o SC special case
- oe or equivalent (and appropriate)
- o dep dependent
- o indep independent
- o awrt answer which rounds to
- o eeoo each error or omission

• No working

If no working is shown then correct answers normally score full marks If no working is shown then incorrect (even though nearly correct) answers score no marks.

• With working

If there is a wrong answer indicated on the answer line always check the working in the body of the script (and on any diagrams), and award any marks appropriate from the mark scheme.

If it is clear from the working that the "correct" answer has been obtained from incorrect working, award 0 marks.

If a candidate misreads a number from the question. Eg. Uses 252 instead of 255; method marks may be awarded provided the question has not been simplified. Examiners should send any instance of a suspected misread to review. If there is a choice of methods shown, mark the method that leads to the answer on the answer line; where no answer is given on the answer line, award the lowest mark from the methods shown.

If there is no answer on the answer line then check the working for an obvious answer.

• Ignoring subsequent work

It is appropriate to ignore subsequent work when the additional work does not change the answer in a way that is inappropriate for the question: eg. Incorrect cancelling of a fraction that would otherwise be correct.

It is not appropriate to ignore subsequent work when the additional work essentially makes the answer incorrect eg algebra.

Transcription errors occur when candidates present a correct answer in working, and write it incorrectly on the answer line; mark the correct answer.

• Parts of questions

Unless allowed by the mark scheme, the marks allocated to one part of the question CANNOT be awarded to another.

International GCSE Maths							
Apart from question 6d, 9b, 15, 18, 21c (where the mark scheme states otherwise) the correct answer, unless clearly obtained from an incorrect method, should be taken to imply a correct method.							
O	Working	Answer	Mark	Notes			
1 a		26 or 64	1	B1 or both 26 and 64 with no others			
b		21 or 39	1	B1 or both 21 and 39 with no others			
с		17 or 31	1	B1 or both 17 and 31 with no others			
d		1 or 64	1	B1 or both 1 and 64 with no others			
				Total 4 marks			

2 a	3000	1	B1
b	6.5	1	B1
			Total 2 marks

3 a		2	M1	for at least 2 correct tallies or
				frequencies
	2, 5, 4, 3, 2		A1	mark frequencies only – in either
				column
b	1	1	B1	allow ft from (a)
c	4	1	B1	
				Total 4 marks

4 a	New York	1	B1 accept -15
b	25	1	B1 accept -25
с	-28	1	B1
			Total 3 marks

5 a		0.003, 0.035, 0.5, 0.539, 0.9	1	B1
b		60	1	B1 allow 60%
с		$8\frac{4}{7}$	1	B1 oe
d	$0.95 - 0.68$ or $\frac{95}{100} - \frac{68}{100}$ or $\frac{19}{20} - 0.68$ oe		2	M1 or $\frac{27}{100}$ or 27%
		0.27		A1
				Total 5 marks

6	а		23	1	B1	
	b		Added 4	1	B1	accept +4, 4 more, oe, $4n - 1$ (need to know 4 and we need to
						add/go up oe)
	С	(23) 27, 31, 35, 39, 43, 47, 51, 55, 59, 63, 67, 71 OR 4 <i>n</i> - 1 = 70		2	M1	allow list of numbers going up in 4's up to 71 or more (allow one error)
			71		A1	
	d	No and identifying all terms in sequence are odd OR No and method to count on as far as 95 (or clearly showing 95) OR No and method to find <i>n</i> when term is 96 e.g. solving $4n - 1 = 96$	No with reason	1	B1	must have 'No' oe or 'is not' oe and a reason.
						Total 5 marks

7 a	(3, -1)	1	B1	
b	(×) at (−2, −4)	1	B1	condone missing label as long on unambiguous
c	(-1, 2)	2	B2	B1 for $(-1, a)$ where $a \neq 2$ or $(b, 2)$ where $b \neq -1$
d	x = 4 drawn	1	B1	
				Total 5 marks

8		3	B3	For the correct time of 13 50 or 1.50 pm or
				1.50 in the afternoon oe
				(B2 for 1.50 or 1.50 am or stating 2 hours 40 mins or 160 mins or intention to add all 4 times onto 11.10
	13 50			B1 for intention to add all 4 times together or evidence of intention to add on 2 or 3 times to 11 10)
				Total 3 marks

9 a i		33	1	B1	accept 32 – 34
ii		15	1	B1	accept 15 – 16
b	e.g. 820 ÷ 10 × "33" (= 2706)		2	M1	method to convert 820 metres to feet or 2850 feet to
	or 2850 ÷ 50 × "15" (= 855)				metres, allow ft from (ai) or (aii) or
					a value for 820 m to feet in range $(2620 - 2740)$ or
					a value for 2850 feet to m in range $(830 - 900)$
		2850 feet		A1	2850 selected (could be unambiguously circled,
		supported by			underlined or stated) with correct working and figures
		working			as above to justify result, ft from part (ai) or (aii)
					Total 4 marks

10	e.g. 360 - (30 + 45 + 165) (= 120)		4	M1	method to calculate One Stop Shoes angle
	e.g. $\frac{30}{45}$, 18 (= 12) or $\frac{30}{120}$, 48 (= 12) oe			M1	method to calculate ABC Shoes frequency
	e.g. $165 \div 45 \times 18 (= 66)$ oe or $165 \div 30 \times "12" (= 66)$ oe or $165 \div "120" \times 48 (= 66)$ oe or $18 + 48$ having shown or implied that $120 + 45 = 165$ and a clear intention that this is the method for Superfast Trainers (= 66)			M1	method to calculate Superfast Trainers frequency
		12, 120, 66		A1	fully correct table
					Total 4 marks

11	0.23 × 450 oe		2	M1 or for an answer of 553.5 or 346.5
		103.5		A1
				Total 2 marks

12	a		1, 3, 9	1	B1	need all three but ignore any repeats
	b	$ \begin{array}{r} 15, 30, 45, 60, 75, 90, 105, 120, 135, 150, 165, 180, 195, \\ 210 \text{ and } 70, 140, 210 \\ \text{OR } 3 \times 5 \text{ and } 2 \times 5 \times 7 \\ \text{OR } 2 \times 3 \times 5 \times 7 (2, 3, 5, 7) \text{ oe eg } \underline{3 \times 5 \times 14} (3, 5, 14) \\ \hline \begin{array}{r} 3 & 15 & 70 \\ \hline 5 & 5 & 70 \\ \hline 7 & 1 & 14 \\ \hline 2 & 1 & 2 \\ \hline & 1 & 1 \\ \end{array} $		2	M1	for listing at least three multiples of 15 and 70 or finding the prime factors of 15 and 70 (could be factors at the ends of branches of factor trees or lists 3, 5 and 2, 5, 7) or a correct calculation or the correct values for the LCM eg 2,3,5,7 or 3,5,14 oe (could be in a table)
			210		A1	
						Total 3 marks

13	eg $\frac{1}{2} \times (20 + 26) \times 12 (= 276)$		5	M2	complete method to find the area of the shape
	or $12 \times 20 + \frac{1}{2} \times (26 - 20) \times 12 (= 276)$ or $12 \times 26 - \frac{1}{2} \times (26 - 20) \times 12 (= 276)$				M1 for method to find the area of a rectangle $12 \times 20 \ (= 240)$ or $12 \times 26 \ (=312)$ or the area of the triangle $\frac{1}{2} \times (26 - 20) \times 12 \ (= 36)$
	"276" ÷ 20 (= 13.8)			M1	(indep) method to find the number of tins for their area ft any value from a calculation that includes at least two of 20, 26 & 12
	eg $3 \times \$40 + 2 \times \13 (= $\$146$) or $14 \times \$13$ (= $\$182$) or $4 \times \$40$ (= $\$160$)			M1	method to calculate a cost for their number of tins dep on previous M1 (NB: use $n \times 40 where <i>n</i> is the next multiple of 4 greater than the number of tins needed, divided by 4)
		146		A1	cao dep on accurate figures
					Total 5 marks

14	а		5(5f-2)	1	B1	
	b	$c+h=5y$ or $\frac{c}{5}=y-\frac{h}{5}$ or $\frac{c+h}{5}$		2	M1	
			$y = \frac{c+h}{5}$		A1	oe if the student puts $\frac{c+h}{5}$ on the answer
						line then if we have previously see
						$y = \frac{c+h}{5}$ we can award full marks
	с	$4x > 2 - 7$ oe or $x + \frac{7}{4} > \frac{2}{4}$ oe		2	M1	accept as an equation or with wrong inequality sign.
			x > -1.25		A1	oe allow $(-1.25, (+)\infty)$
						Note: award M1A0 for an answer of
						-1.25 with no sign or the incorrect sign
						eg $x = -1.25, x < -1.25$
						Total 5 marks

15	$\frac{2}{5} \times \frac{20}{11}$ or $eg \frac{8}{20} \div \frac{11}{20}$		2	M1	For inverting $\frac{11}{20}$ and a clear intention to multiply or for writing both fractions correctly over the same common denominator
	$\frac{\frac{2}{5} \times \frac{20}{11} = \frac{40}{55} = \frac{8}{11} \text{ or}}{\frac{2}{5}^{1} \times \frac{20}{11} = \frac{8}{11}} \text{ or}}$ $\frac{\frac{8}{20} \div \frac{11}{20} = \frac{8}{11}}{\frac{8}{11}}$	Clearly shown		A1	dep on M1 continued to clearly show given result
					Total 2 marks

16	а		$50 < L \le 60$	1	B1	oe eg 50 - 60
	b	25 × 6 + 35 × 26 + 45 × 31 + 55 × 40 + 65 × 17 (150 + 910 + 1395 + 2200 + 1105)(= 5760)			M2	For correct products using midpoints (allow one error) with intention to add. M1 for products using frequency and a consistent value within the range (allow one error) with intention to add or correct products using midpoints
						(allow one error) without addition
		"5760" ÷ "120"			M1	dep on M1
			48	4	A1	
						Total 5 marks

17 a	$150 \div 6 \times 14$ oe		2	M1
		350		A1
b	$630 \div 90 \times 6$ oe		2	M1
		42		A1
с	$162 \div (2+7) \times 2$ oe		2	M1
		36		A1
				Total 6 marks

18	<i>ADC</i> = 180 – 58 (= 122) or <i>EDF</i> = 122 or <i>CDE</i> = 58 or <i>ADF</i> = 58		5	M1	may be seen marked on the diagram
	e.g. $DEF = 58 \div 2$ or $DEF = (180 - 122) \div 2$			M1	complete method to find angle <i>DEF</i>
		29		A1	
				B2	dep on M2 for fully correct reasons for their method (B1 dep on M1 for one correct reason stated and used) e.g. <u>Allied angles, co-interior angles, Alternate angles,</u> <u>Corresponding angles, Vertically opposite angles are</u> equal (or Vertically <u>opposite angles</u> are equal), <u>Angles</u> on a straight <u>line</u> add up to 180°(or angles on a straight <u>line</u> add to <u>180°</u>), Sum of <u>two angles</u> in a triangle are equal to <u>opposite exterior</u> angle, <u>Angles</u> in a <u>triangle</u> add up to 180°(or Angles in a <u>triangle</u> add up to <u>180°</u>), Base angles in an <u>isosceles</u> triangle <u>Angles</u> in a <u>quadrilateral</u> add up to 360. (accept "4-sided shape" or parallelogram) <u>Opposite angles</u> of a <u>parallelogram</u> are equal
					Total 5 marks

19	$76 \div (5 + 2 - 3) (= 19)$		4	M1
	$5x + 2x - 3x = 76$, $x = 76 \div 4$ (=19) oe			
	3 × "19" (= 57)			M1
	"57" – 48.5(0)			M1
		8.5(0)		A1
				Total 4 marks

20	а	1.04 × 3 130 000 oe		3	M2	complete method to incr	cease salary by 4%		
							M1 for 0.04 × 3 130 000 (= 125 200)) oe	
				3 255 200		A1			
	b	for 0.15 × 750 000 oe (=112 500) or 0.85 × 750 000 oe (637 500)	OR		3	M1	For method to find depreciation for 1 year or value after 1 year	OR M2 for 750 000 × 0.85 ³ (= 460 593.75) or 750 000 × 0.85 ⁴	
		0.85 × "637 500" (= 541 875) oe 0.85 × "541 875" (= 460 593.75) oe	$\begin{array}{ c c c c }\hline 750\ 000 \\ \times\ 0.85^3 \end{array}$			M1	for completing method	(= 391 504.69) (M1 for 750 000 × 0.85 ² (= 541 875)	
				460 594		A1	accept 460 593 - 460 594		
							SC: if no other marks gained award M1 for 0.55×750000 oe (= 412 500) or 0.45×750000 oe (= 337 500) accept (1 – 0.15) as equivalent to 0.85 throughout		
								Total 6 marks	

21	а			g^{10}	1	B1	
	b			$9c^2d^8$	2	B2	B1 for 2 out of 3 terms correct as part of a product
	с	eg $4x + 3y = 17$ - 4x + 8y = 20 or eg $4(5 - 2y) + 3y = 17$	eg $8x + 6y = 34$ - 3x + 6y = 15 or eg $4x + 3 \times \frac{1}{2}(5 - x) = 17$		3	M1	Correct method to eliminate x or y: coefficients of x or y the same and correct operation to eliminate selected variable (condone any one arithmetic error in multiplication) or writing x or y in terms of the other variable and correctly substituting
		eg $4x + 3 \times 0.6 = 17$ or $x + 2 \times 0.6 = 5$	eg $4 \times 3.8 + 3y = 17$ or $3.8 + 2y = 5$	r - 3.8		M1	(dep) correct method to find second variable – could start process again or use substitution
				x = 5.8 y = 0.6		AI	M1
							Total 6 marks

22	$\tan x = \frac{3.4}{4.7}$ oe eg $\cos x = \frac{4.7}{\sqrt{3.4^2 + 4.7^2}}$			M1	or $\sin x = \frac{3.4 \sin 90}{\sqrt{3.4^2 + 4.7^2}}$ oe
	$(x =) \tan^{-1}\left(\frac{3.4}{4.7}\right)$ oe eg $(x =) \cos^{-1}\left(\frac{4.7}{\sqrt{3.4^2 + 4.7^2}}\right)$			M1	or $(x =)\sin^{-1}\left(\frac{3.4\sin 90}{\sqrt{3.4^2 + 4.7^2}}\right)$ oe
		35.9	3	A1	accept 35.7 - 36.1
					Total 3 marks

PMT

23	$8.5^2 - (8 \div 2)^2 (= 56.25) \text{ or } \cos x = \frac{4}{8.5}$		4	M1	or eg cos $A = \frac{8^2 + 8.5^2 - 8.5^2}{2 \times 8 \times 8.5}$
	$\sqrt{56.25}$ (= 7.5) or $x = \cos^{-1}\left(\frac{4}{8.5}\right)$ (= 61.927)			M1	or eg $(A =)\cos^{-1}\left(\frac{8^2 + 8.5^2 - 8.5^2}{2 \times 8 \times 8.5}\right)$ (61.927) (other angle = 56.144)
	$8 \times "7.5" \div 2$ oe or $0.5 \times 8 \times 8.5 \times \sin "61.927"$			M1	or eg 0.5 × 8.5 × 8 × sin61.927oe
		30		A1	
					Total 4 marks

24	$\pi \times 3^2 \times h = 72\pi$ oe		5	M1	Allow use of 3.14 or $\frac{22}{7}$ for π and use of 226 for 72π
	$h = 72\pi \div (\pi \times 3^2)$ oe or $h = 8$			M1	method to isolate h (may be seen in several stages)
	$2 \times \pi \times 3^2$ (= 18 π or 56.54) or $2 \times \pi \times 3 \times$ "8" oe (= 48 π or 150 - 151)			M1	method to find the area of the two circles or curved surface area – use of their <i>h</i> , dep on M1 (NB may get this mark for area of 2 circles with no previous marks awarded)
	$2 \times \pi \times 3^2 + 2 \times \pi \times 3 \times \text{``8'' oe} (= 66\pi)$	207		M1	complete method to find the total surface area ft their h dep on 1st M1, including intention to add, to find the total surface area
		207		A1	accept 207-208
					Total 5 marks

PMT

Pearson Education Limited. Registered company number 872828 with its registered office at 80 Strand, London, WC2R 0RL, United Kingdom