

CAMBRIDGE INTERNATIONAL EXAMINATIONS

Cambridge International General Certificate of Secondary Education

MARK SCHEME for the March 2016 series**0606 ADDITIONAL MATHEMATICS****0606/12**

Paper 12, maximum raw mark 80

This mark scheme is published as an aid to teachers and candidates, to indicate the requirements of the examination. It shows the basis on which Examiners were instructed to award marks. It does not indicate the details of the discussions that took place at an Examiners' meeting before marking began, which would have considered the acceptability of alternative answers.

Mark schemes should be read in conjunction with the question paper and the Principal Examiner Report for Teachers.

Cambridge will not enter into discussions about these mark schemes.

Cambridge is publishing the mark schemes for the March 2016 series for most Cambridge IGCSE® and Cambridge International A and AS Level components.

® IGCSE is the registered trademark of Cambridge International Examinations.

Page 2	Mark Scheme	Syllabus	Paper
	Cambridge IGCSE – March 2016	0606	12

Abbreviations

awrt	answers which round to
cao	correct answer only
dep	dependent
FT	follow through after error
isw	ignore subsequent working
oe	or equivalent
rot	rounded or truncated
SC	Special Case
soi	seen or implied
www	without wrong working

Question	Answer	Marks	Guidance
1	$ax + 9 = -2x^2 + 3x + 1$ $2x^2 + (a - 3)x + 8 = 0$ For 2 distinct roots, $(a - 3)^2 > 64$ Critical values -5 and 11 $a > 11$, $a < -5$	M1 M1 A1 A1	for attempt to equate the line and the curve and obtain a 3 term quadratic equation for use of the discriminant for critical values for correct range
2	$a = -\frac{13}{6}$, $b = 0$, $c = 1$	B3	B1 for each
3	$\log_5 \sqrt{x} + \log_{25} x = 3$ $\frac{1}{2} \log_5 x + \frac{\log_5 x}{\log_5 25} = 3$ $\log_5 x = 3$ $x = 125$ cao Alternative scheme: $\frac{\log_{25} \sqrt{x}}{\log_{25} 5} + \log_{25} x = 3$ $\frac{1}{2} \log_{25} x + \log_{25} x = 3$ $\log_{25} x = \frac{3}{2}$ $x = 125$ cao	B1,B1 B1 B1 B1	B1 for $\frac{1}{2} \log_5 x$ B1 for $\frac{\log_5 x}{\log_5 25}$ for final answer for change of base for $\frac{1}{2} \log_{25} x$ (must be from correct work) for final answer

Page 3	Mark Scheme	Syllabus	Paper
	Cambridge IGCSE – March 2016	0606	12

Question	Answer	Marks	Guidance
4 (i)		B1 B1 B1 B1	for a line in correct position for (0, 2), (2, 0) for correct shape for $y = 3 + 2x $, touching the x -axis for (-1.5, 0), (0, 3)
(ii)	$2 - x = 3 + 2x$ leading to $x = -\frac{1}{3}$ $2 - x = -3 - 2x$ leading to $x = -5$ Alternative: $(2 - x)^2 = (3 + 4x)^2$ leading to $15x^2 + 28x + 5 = 0$ $x = -\frac{1}{3}, x = -5$	B1 M1 A1 M1 A1,A1	for $x = -\frac{1}{3}$ for correct attempt to deal with 'negative' branch. for $x = -5$ for equating and squaring to obtain a 3 term quadratic equation A1 for each.
5 (a) (i)	${}^9P_6 = 60480$	B1	Must be evaluated
(ii)	${}^4P_2 \times {}^3P_2 \times 2 = 144$	M1,A1	M1 for attempt a product of 3 perms
(iii)	840×2 1680	B1,B1	B1 for either 840, or realising that there are 2 possible positions for the symbols
(b) (i)	${}^{10}C_6 \times {}^5C_3$ 2100	M1 A1	for unsimplified form
(ii)	${}^8C_4 \times {}^4C_2$ 420	M1 A1	for unsimplified form
6 (i)	$f(x) > 6$	B1	Allow B1 for $y > 6$
(ii)	$f^{-1}(x) = \frac{1}{4} \ln(x - 6)$ Domain: $x > 6$ Range: $f^{-1}(x) \in \mathbb{R}$	M1 A1 B1 B1	for a complete method must be $f^{-1}(x) =$ or $y = \dots$ must be using the correct variable in both
(iii)	$f'(x) = 4e^{4x}$	B1	
(iv)	$6 + e^{4x} = 4e^{4x}$ leading to $x = \frac{1}{4} \ln 2$	M1 A1	for a complete, correct method

Page 4	Mark Scheme	Syllabus	Paper
	Cambridge IGCSE – March 2016	0606	12

Question	Answer	Marks	Guidance
7 (i)	$f\left(\frac{1}{2}\right) = \frac{a}{8} + \frac{7}{4} - \frac{9}{2} + b \quad (=0)$ $a + 8b = 22$ $8a + 28 - 18 + b = 5(-a + 7 + 9 + b)$ $13a - 4b = 70$ <p>leading to $a = 6, b = 2$</p>	M1	for attempt at $f\left(\frac{1}{2}\right)$
(ii)	$(2x-1)(3x^2 + 5x - 2)$	B2,1,0	-1 each error
(iii)	$(2x-1)(3x-1)(x+2)$	M1 A1FT	for attempt to factorise their quadratic factor must be 3 linear factors
8 (i)	$\lg y = \lg A + b \lg x$ Gradient = 1.2 so $b = 1.2$ Intercept = 1.44 $A = 27.5$	B1 M1 A1	may be implied by later work for attempt at gradient for $b = 1.2$
(ii)	when $x = 100, \lg x = 2$ $\lg y = 3.84$ (allow 3.8 to 3.9)	M1 A1	for correct use of graph or equation
(iii)	when $y = 8000, \lg 8000 = 3.9, \lg x = 2.05$ leading to $x = 113, 10^{2.05}$ or 112	M1 A1	for correct use of graph or equation

Page 5	Mark Scheme	Syllabus	Paper
	Cambridge IGCSE – March 2016	0606	12

Question	Answer	Marks	Guidance	
9 (i)	$\frac{7}{2}r^2\theta = \frac{1}{2}r^2(2\pi - \theta)$ $\theta = \frac{\pi}{4} \text{ oe}$	M1	for a valid method	
		A1	allow in degrees	
	(ii)	$r + r + \frac{\pi}{4}r = 20$, leading to $r = 7.180(3..)$	M1	for valid method
			A1	Must show enough accuracy to get A1
(iii)	Perimeter = $\frac{\pi}{4}r + 2r \tan \frac{\pi}{8}$ $= 5.6394 + 5.9484$ $= 11.6$	B1,B1	B1 for arc length, B1 for twice AC	
		B1	for 11.6	
(iv)	Area = $(r \times AC) - \frac{1}{2}r^2 \frac{\pi}{4}$ $= 21.356 - 20.246$ or equivalent method using triangles	B1,B1	B1 for area of quadrilateral, allow unsimplified, B1 for sector area	
		B1	for area in given range	
10 (i)	$x \times \frac{3}{2} \times 2(2x-1)^{\frac{1}{2}} + (2x-1)^{\frac{3}{2}}$	B1	for $\frac{3}{2} \times 2(2x-1)^{\frac{1}{2}}$	
		M1	for attempt at differentiation of a product	
		A1	for all else correct	
(ii)	$3 \int x(2x-1)^{\frac{1}{2}} dx = x(2x-1)^{\frac{3}{2}} - \int (2x-1)^{\frac{3}{2}} dx$ $= x(2x-1)^{\frac{3}{2}} - \frac{1}{2} \times \frac{2}{5}(2x-1)^{\frac{5}{2}}$	M1	for attempt to use part (i)	
		B1,B1	B1 for $x(2x-1)^{\frac{3}{2}}$, allow if divided by 3 B1 for $\frac{1}{2} \times \frac{2}{5}(2x-1)^{\frac{5}{2}}$, allow if divided by 3	
(iii)	$\int x(2x-1)^{\frac{1}{2}} dx = \frac{1}{3}(2x-1)^{\frac{3}{2}} \left(x - \frac{1}{5}(2x-1) \right)$ $= \frac{(2x-1)^{\frac{3}{2}}}{15} (3x+1)$	M1	for taking out a common factor of $(2x-1)^{\frac{3}{2}}$	
		DM1	for attempt to obtain a linear factor	
		A1		
(iii)	$\left(\frac{1}{15} \times 4 \right) - 0$	M1	for attempt to use limits correctly	
		A1FT	FT on their $\frac{px+q}{15}$	

Page 6	Mark Scheme	Syllabus	Paper
	Cambridge IGCSE – March 2016	0606	12

Question	Answer	Marks	Guidance			
11 (i)	$\frac{1}{\operatorname{cosec}\theta - 1} - \frac{1}{\operatorname{cosec}\theta + 1} = \frac{\operatorname{cosec}\theta + 1 - \operatorname{cosec}\theta + 1}{\operatorname{cosec}^2\theta - 1}$ $= \frac{2}{\cot^2\theta}$ $= 2 \tan^2\theta$	M1	for attempt to obtain a single fraction			
		A1	all correct as shown			
		M1	for use of correct identity			
		A1	for 'finishing off'			
	Alternative scheme:	$\frac{1}{\operatorname{cosec}\theta - 1} - \frac{1}{\operatorname{cosec}\theta + 1} = \frac{\sin\theta}{1 - \sin\theta} - \frac{\sin\theta}{1 + \cos\theta}$ $= \frac{(\sin\theta + \sin^2\theta) - (\sin\theta - \sin^2\theta)}{1 - \sin^2\theta}$ $= \frac{2\sin^2\theta}{\cos^2\theta}$ $= 2 \tan^2\theta$	M1	for attempt to obtain a single fraction in terms of $\sin\theta$ only		
			A1	all correct as shown		
			M1	for use of correct identity		
			A1	for 'finishing off'		
			(ii)	$2 \tan^2\theta = 6 + \tan\theta$ $(2 \tan\theta + 3)(\tan\theta - 2) = 0$ $\tan\theta = -\frac{3}{2}, \tan\theta = 2$	M1	for attempt to use (i), to obtain a quadratic equation and valid attempt to solve
					DM1	for attempt to solve trig equation
	$\theta = 63.4^\circ, 123.7^\circ, 243.4^\circ, 303.7^\circ$	A1,A1	for each 'pair'			