

**UNIVERSITY OF CAMBRIDGE INTERNATIONAL EXAMINATIONS**  
International General Certificate of Secondary Education

**MARK SCHEME for the October/November 2010 question paper  
for the guidance of teachers**

**0580 MATHEMATICS**

**0580/43**

Paper 4 (Extended), maximum raw mark 130

This mark scheme is published as an aid to teachers and candidates, to indicate the requirements of the examination. It shows the basis on which Examiners were instructed to award marks. It does not indicate the details of the discussions that took place at an Examiners' meeting before marking began, which would have considered the acceptability of alternative answers.

Mark schemes must be read in conjunction with the question papers and the report on the examination.

- CIE will not enter into discussions or correspondence in connection with these mark schemes.

CIE is publishing the mark schemes for the October/November 2010 question papers for most IGCSE, GCE Advanced Level and Advanced Subsidiary Level syllabuses and some Ordinary Level syllabuses.



<b>Page 2</b>	<b>Mark Scheme: Teachers' version</b>	<b>Syllabus</b>	<b>Paper</b>
	<b>IGCSE – October/November 2010</b>	<b>0580</b>	<b>43</b>

### Abbreviations

cao	correct answer only
cso	correct solution only
dep	dependent
ft	follow through after error
isw	ignore subsequent working
oe	or equivalent
SC	Special Case
www	without wrong working
art	anything rounding to
soi	seen or implied

Qu.	Answers	Mark	Part Marks
<b>1</b>	<p><b>(a)</b> <math>200 \div 10 \times 3</math> oe <math>200 \div 10 \times 2</math> oe</p> <p><b>(b)</b> 65</p> <p><b>(c)</b> 46</p> <p><b>(d)</b> 0.6(0)</p>	<p>M1 M1</p> <p>2</p> <p>3</p> <p>3</p>	<p><b>M1</b> for <math>\frac{39}{60} \times 100</math> oe 35 is <b>M0</b></p> <p><b>M2</b> for <math>36.80 \div 0.8</math> oe or <b>M1</b> for <math>80\% = 36.80</math> oe</p> <p><b>M2</b> for <math>5(x + 12) + 2x = 64.2</math> oe or <math>(64.2 - 5 \times 12) \div 7</math> or <math>5x + 2(x - 12) = 64.2</math> oe or <math>(64.2 + 2 \times 12) \div 7</math> or <b>M1</b> for <math>y = x + 12</math> and <math>5y + 2x = 64.2</math> or <math>y = x - 12</math> and <math>5x + 2y = 64.2</math> After <b>M0</b>, <b>SC1</b> for <math>k(x \pm 12)</math> seen</p>
<b>2</b>	<p><b>(a)</b> <math>(\cos Q =) \frac{4^2 + 4.5^2 - 7^2}{2 \times 4 \times 4.5}</math> o.e. 110.74....</p> <p><b>(b)</b> <math>(RS =) \frac{7 \sin 40}{\sin 85}</math> 4.516 ...</p> <p><b>(c)</b> Angle <math>R = 55^\circ</math> <math>0.5 \times 7 \times 4.52 \times \sin(\text{their } 55)</math> o.e. <math>0.5 \times 4 \times 4.5 \times \sin 110.7</math> o.e. Triangle <math>PRS</math> + Triangle <math>PQR</math> 21.4 (21.36 – 21.42)</p>	<p>M2 E2</p> <p>M2 E1</p> <p>B1 M1 M1 M1 A1</p>	<p><b>M1</b> for <math>7^2 = 4^2 + 4.5^2 - 2 \times 4 \times 4.5 \times \cos(Q)</math></p> <p>If <b>E0</b> then <b>A1</b> for <math>-0.354(1\dots)</math></p> <p><b>M1</b> for <math>\frac{RS}{\sin 40} = \frac{7}{\sin 85}</math> o.e.</p> <p>Can be implied by second M</p> <p>(May be seen on diagram)</p> <p>(12.95 – 13.0) their 55 is (180 – 40 – 85)</p> <p>(8.418 – 8.42) (<math>s = 7.75</math>)</p> <p>Dependent on M1, M1</p> <p>www 5</p>

Page 3	Mark Scheme: Teachers' version	Syllabus	Paper
	IGCSE – October/November 2010	0580	43

3	<p>(a) <math>5x^2 - x</math> or <math>x(5x - 1)</math></p> <p>(b) <math>27x^9</math></p> <p>(c) (i) <math>7x^7(1 + 2x^7)</math></p> <p>(ii) <math>(y + w)(x + 2a)</math></p> <p>(iii) <math>(2x + 7)(2x - 7)</math></p> <p>(d) <math>\frac{-5 \pm \sqrt{5^2 - 4(2)(1)}}{2(2)}</math> oe</p> <p>-2.28 -0.22</p>	2 2 2 2 1 2 1 1	<p><b>M1</b> for <math>x^2 + 3x</math> or <math>4x^2 - 4x</math> correct</p> <p><b>B1</b> for 27 or for <math>x^9</math></p> <p><b>M1</b> for any correct partially factorised expression or <math>7x^7(1 + \dots)</math></p> <p><b>M1</b> for <math>x(y + w) + 2a(y + w)</math> or <math>y(x + 2a) + w(x + 2a)</math></p> <p>In square root <b>B1</b> for <math>5^2 - 4(2)(1)</math> or better (17)</p> <p>If in form <math>\frac{p + \sqrt{q}}{r}</math> or <math>\frac{p - \sqrt{q}}{r}</math></p> <p><b>B1</b> for <math>p = -5</math> and <math>r = 2(2)</math></p> <p><b>SC1</b> for -2.3 or -2.281 to -2.280 and -0.2 or -0.220 to -0.219</p>
4	<p>(a) (i) <math>\begin{pmatrix} 25 \\ 43 \end{pmatrix}</math></p> <p>(ii) (16)</p> <p>(iii) <math>\frac{1}{-2} \begin{pmatrix} 5 &amp; -3 \\ -4 &amp; 2 \end{pmatrix}</math> isw or <math>\begin{pmatrix} \frac{5}{2} &amp; \frac{3}{2} \\ 2 &amp; -1 \end{pmatrix}</math></p> <p>(b) Reflection only x-axis oe</p> <p>(c) <math>\begin{pmatrix} 0 &amp; -1 \\ 1 &amp; 0 \end{pmatrix}</math></p>	1 1 2 2 1 1 2	<p>If 0, 0 then <b>SC1</b> for 25 and 43 seen</p> <p><b>B1</b> for 16 without brackets</p> <p><b>B1</b> for determinant = -2</p> <p>or <b>B1</b> for <math>k \begin{pmatrix} 5 &amp; -3 \\ -4 &amp; 2 \end{pmatrix}</math></p> <p>If more than one transformation given – no marks available</p> <p>independent</p> <p><b>B1</b> for one correct column</p>

Page 4	Mark Scheme: Teachers' version	Syllabus	Paper
	IGCSE – October/November 2010	0580	43

5	(a) (i) Accurate perpendicular bisector, with 2 pairs of arcs, of $CD$ .	2	SC1 if accurate without arcs.	
	(ii) Accurate angle bisector, with two pairs of arcs, of angle $A$ .	2	SC1 if accurate without arcs.	
	(b) SHOP written in correct region	S1	Dependent on at least SC1 in (i) and (ii) and intersection	
	(c) (i) Arc, centre $B$ , radius 5cm, reaching across $ABCD$ .	1	Allow good freehand	
	(ii) Area outside their arc centre $B$ and outside SHOP shaded	1ft	dep on S1	
6	(a) (i) 33	1	<b>Accept</b> fraction, %, dec equivalents (3sf or better) throughout but not ratio or words i.s.w. incorrect cancelling/conversion to other forms Pen –1 once for 2 sf answers	
	(ii) $\frac{243}{3125}$ (0.07776)	2		Accept 0.0778. <b>M1</b> for $\left(\frac{3}{5}\right)^5$ oe
	(b) (i) $\frac{2}{5}, \frac{3}{4}, \frac{1}{8}, \frac{7}{8}$	3		<b>B1</b> for $\frac{2}{5}$ and $\frac{3}{4}$ <b>B1</b> for $\frac{1}{8}$ <b>B1</b> for $\frac{7}{8}$
	(ii) $\frac{1}{20}$ (0.05) cao	2		<b>M1</b> for their $\frac{2}{5} \times$ their $\frac{1}{8}$
	(iii) $\frac{1}{5}$ (0.2) ft	2ft		ft $\frac{3}{20}$ + their (b)(ii) or <b>M1</b> for $\frac{3}{5} \times \frac{1}{4}$
7	(a) – 5.4 3.7	1 1	<b>P3ft</b> their table. <b>P2ft</b> for 6 or 7 points. <b>P1ft</b> for 4 or 5 points Only ft points if shape not affected.	
	(b) 8 points correctly plotted ft  Smooth cubic curve through all 8 points	P3  C1		
	(c) –2, –4, 4	2		<b>B1</b> for 2 correct
	(d) 7 points correctly plotted ft Two <b>separate</b> smooth branches of rectangular hyperbola	P2 C1		<b>P2ft</b> <b>P1ft</b> for 5 or 6 points Must pass through all 7 points, only ft if shape not affected and no contact with either axis.
	(e) (i) $-2.9 \leq x \leq -2.8$ $2.05 \leq x \leq 2.15$	1 1		Not with $y$ coordinates
	(ii) $a = 10$	1		
	$b = -40$	1		

Page 5	Mark Scheme: Teachers' version	Syllabus	Paper
	IGCSE – October/November 2010	0580	43

8	<p>(a) (i) 396 (395.6 – 396)</p> <p>(ii) 3.13 (3.125 – 3.128....) ft</p> <p>(iii) 144 (144 – 144.4) ft</p> <p>(b) (i) 311 (310.8 – 311.1)</p> <p>(ii) 3.50 (3.496 to 3.50) ft</p>	<p>4</p> <p>2ft</p> <p>2ft</p> <p>5</p> <p>2ft</p>	<p><b>M1</b> for <math>\frac{2}{3} \times \pi \times 3^3</math> and <b>M1</b> (independent) for <math>\pi \times 3^2 \times 12</math>,</p> <p><b>M1</b> (dependent on M2) for adding 126 <math>\pi</math> implies <b>M3</b></p> <p>ft their (i) <math>\times 7.9 \div 1000</math>.</p> <p><b>M1</b> for <math>\times 7.9</math> soi by figs 313 or 3125 – 3128...</p> <p>ft <math>15 \times 6 \times 6</math> – their (a)(i)</p> <p><b>M1</b> for <math>6 \times 6 \times 15</math> oe</p> <p><b>M1</b> for <math>2 \times \pi \times 3^2</math> and <b>M1</b> (independent) for <math>\pi \times 6 \times 12</math> and <b>M1</b> for <math>\pi \times 3^2</math>,</p> <p><b>M1</b> (dependent on M3) for adding. (99<math>\pi</math> implies <b>M4</b>)</p> <p>ft their (b)(i) <math>\times 0.01125</math></p> <p><b>M1</b> for their (b)(i) <math>\div 8</math> and <math>\times</math> figs 9 implied by figs 3496 to 350</p>
9	<p>(a) (i) <math>\begin{pmatrix} 9 \\ 5 \end{pmatrix}</math></p> <p>(ii) <math>\begin{pmatrix} 4 \\ 7 \end{pmatrix}</math></p> <p>(iii) <math>\overrightarrow{BA}</math> or <math>-\overrightarrow{AB}</math></p> <p>(iv) 10.3 (10.29 – 10.30)</p> <p>(b) (i) <math>2\mathbf{u}</math></p> <p>(ii) <math>\frac{1}{2}(\mathbf{t} - \mathbf{u})</math> oe</p> <p>(iii) <math>\frac{3}{2}\mathbf{u} + \frac{1}{2}\mathbf{t}</math> oe ft</p>	<p>1</p> <p>1</p> <p>1</p> <p>1</p> <p>2</p> <p>1</p> <p>2</p> <p>2ft</p>	<p>If 0, <b>SC1</b> for <math>\overrightarrow{CB} = \begin{pmatrix} 5 \\ -2 \end{pmatrix}</math> seen</p> <p><math>BA</math> not indicated as a vector is not enough.</p> <p><b>M1</b> for (their 9)<sup>2</sup> + (their 5)<sup>2</sup></p> <p><b>M1</b> for <math>\frac{1}{2}</math> (their <math>\overrightarrow{BA} + \overrightarrow{AD} + \overrightarrow{DC}</math>) or equivalent</p> <p>correct route for <math>\overrightarrow{BM}</math>, along obtainable vectors in terms of <math>\mathbf{t}</math> and <math>\mathbf{u}</math></p> <p>or <b>M1</b> for correct unsimplified answer</p> <p>ft their (i) + their (ii) simplified</p> <p>or <math>\mathbf{t} + \mathbf{u}</math> – their (b)(ii) simplified</p> <p><b>M1</b> for correct (or ft) unsimplified (i) + (ii)</p> <p>or <math>\mathbf{t} + \mathbf{u}</math> – their (b)(ii)</p>

