



Cambridge IGCSE™

CANDIDATE
NAME

CENTRE
NUMBER

--	--	--	--	--

CANDIDATE
NUMBER

--	--	--	--



ADDITIONAL MATHEMATICS

0606/22

Paper 2

October/November 2021

2 hours

You must answer on the question paper.

No additional materials are needed.

INSTRUCTIONS

- Answer **all** questions.
- Use a black or dark blue pen. You may use an HB pencil for any diagrams or graphs.
- Write your name, centre number and candidate number in the boxes at the top of the page.
- Write your answer to each question in the space provided.
- Do **not** use an erasable pen or correction fluid.
- Do **not** write on any bar codes.
- You should use a calculator where appropriate.
- You must show all necessary working clearly; no marks will be given for unsupported answers from a calculator.
- Give non-exact numerical answers correct to 3 significant figures, or 1 decimal place for angles in degrees, unless a different level of accuracy is specified in the question.

INFORMATION

- The total mark for this paper is 80.
- The number of marks for each question or part question is shown in brackets [].

This document has **16** pages. Any blank pages are indicated.

Mathematical Formulae**1. ALGEBRA***Quadratic Equation*

For the equation $ax^2 + bx + c = 0$,

$$x = \frac{-b \pm \sqrt{b^2 - 4ac}}{2a}$$

Binomial Theorem

$$(a + b)^n = a^n + \binom{n}{1}a^{n-1}b + \binom{n}{2}a^{n-2}b^2 + \dots + \binom{n}{r}a^{n-r}b^r + \dots + b^n$$

where n is a positive integer and $\binom{n}{r} = \frac{n!}{(n-r)!r!}$

Arithmetic series $u_n = a + (n-1)d$

$$S_n = \frac{1}{2}n(a+l) = \frac{1}{2}n\{2a + (n-1)d\}$$

Geometric series $u_n = ar^{n-1}$

$$S_n = \frac{a(1-r^n)}{1-r} \quad (r \neq 1)$$

$$S_\infty = \frac{a}{1-r} \quad (|r| < 1)$$

2. TRIGONOMETRY*Identities*


$$\begin{aligned} \sin^2 A + \cos^2 A &= 1 \\ \sec^2 A &= 1 + \tan^2 A \\ \operatorname{cosec}^2 A &= 1 + \cot^2 A \end{aligned}$$

Formulae for $\triangle ABC$

$$\begin{aligned} \frac{a}{\sin A} &= \frac{b}{\sin B} = \frac{c}{\sin C} \\ a^2 &= b^2 + c^2 - 2bc \cos A \\ \Delta &= \frac{1}{2}bc \sin A \end{aligned}$$

3

1



- (a) On the axes, draw the graphs of $y = 5 + |3x - 2|$ and $y = 11 - x$. [4]
- (b) Using the graphs, or otherwise, solve the inequality $11 - x < 5 + |3x - 2|$. [2]

2 (a) Expand $(2 - 3x)^4$, evaluating all of the coefficients. [4]

(b) The sum of the first three terms in ascending powers of x in the expansion of $(2 - 3x)^4 \left(1 + \frac{a}{x}\right)$ is $\frac{32}{x} + b + cx$, where a , b and c are integers. Find the values of each of a , b and c . [4]

3 (a) Show that $\frac{1}{\sec x - 1} + \frac{1}{\sec x + 1} = 2 \cot x \operatorname{cosec} x$. [4]

(b) Hence solve the equation $\frac{1}{\sec x - 1} + \frac{1}{\sec x + 1} = 3 \sec x$ for $0^\circ < x < 360^\circ$. [4]

6

- 4 (a) Find the x -coordinates of the stationary points on the curve $y = 3 \ln x + x^2 - 7x$, where $x > 0$. [5]

- (b) Determine the nature of each of these stationary points. [3]

7


5 (a) Solve the following simultaneous equations.

$$\begin{aligned}e^x + e^y &= 5 \\ 2e^x - 3e^y &= 8\end{aligned}\quad [5]$$

(b) Solve the equation $e^{(2t-1)} = 5e^{(5t-3)}$. [4]

6 DO NOT USE A CALCULATOR IN THIS QUESTION.

All lengths in this question are in centimetres.



You may use the following trigonometrical ratios.

$$\sin 60^\circ = \frac{\sqrt{3}}{2}$$

$$\cos 60^\circ = \frac{1}{2}$$

$$\tan 60^\circ = \sqrt{3}$$

The diagram shows triangle ABC with $AC = \sqrt{6} - \sqrt{2}$, $AB = \sqrt{6} + \sqrt{2}$ and angle $CAB = 60^\circ$.

(a) Find the exact length of BC .

[3]

(b) Show that $\sin ACB = \frac{\sqrt{6} + \sqrt{2}}{4}$.

[2]

(c) Show that the perpendicular distance from A to the line BC is 1.

[2]

7 It is given that $\frac{d^2y}{dx^2} = e^{2x} + \frac{1}{(x+1)^2}$ for $x > -1$.

(a) Find an expression for $\frac{dy}{dx}$ given that $\frac{dy}{dx} = 2$ when $x = 0$. [3]

(b) Find an expression for y given that $y = 4$ when $x = 0$. [3]

10

8 Variables x and y are such that when \sqrt{y} is plotted against $\log_2(x+1)$, where $x > -1$, a straight line is obtained which passes through $(2, 10.4)$ and $(4, 15.4)$.

(a) Find \sqrt{y} in terms of $\log_2(x+1)$. [4]

(b) Find the value of y when $x = 15$. [1]

(c) Find the value of x when $y = 25$.

[3]

- 9 (a) Find the equation of the normal to the curve $y = x^3 + x^2 - 4x + 6$ at the point (1, 4). [5]

(b) **DO NOT USE A CALCULATOR IN THIS PART OF THE QUESTION.**

Find the exact x -coordinate of each of the two points where the normal cuts the curve again. [5]

- 10 (a)** The first three terms of an arithmetic progression are x , $5x - 4$ and $8x + 2$. Find x and the common difference. [4]

15

(b) The first three terms of a geometric progression are y , $5y - 4$ and $8y + 2$.

(i) Find the two possible values of y .

[4]

(ii) For each of these values of y , find the corresponding value of the common ratio.

[2]

BLANK PAGE

Permission to reproduce items where third-party owned material protected by copyright is included has been sought and cleared where possible. Every reasonable effort has been made by the publisher (UCLES) to trace copyright holders, but if any items requiring clearance have unwittingly been included, the publisher will be pleased to make amends at the earliest possible opportunity.

To avoid the issue of disclosure of answer-related information to candidates, all copyright acknowledgements are reproduced online in the Cambridge Assessment International Education Copyright Acknowledgements Booklet. This is produced for each series of examinations and is freely available to download at www.cambridgeinternational.org after the live examination series.

Cambridge Assessment International Education is part of the Cambridge Assessment Group. Cambridge Assessment is the brand name of the University of Cambridge Local Examinations Syndicate (UCLES), which itself is a department of the University of Cambridge.