

Cambridge IGCSE®

CANDIDATE
NAME

--	--	--	--	--

CENTRE
NUMBER

--	--	--	--	--

CANDIDATE
NUMBER

--	--	--	--



ADDITIONAL MATHEMATICS

0606/02

Paper 2

For examination from 2020

SPECIMEN PAPER

2 hours

You must answer on the question paper.

No additional materials are needed.

INSTRUCTIONS

- Answer **all** questions.
- Use a black or dark blue pen. You may use an HB pencil for any diagrams or graphs.
- Write your name, centre number and candidate number in the boxes at the top of the page.
- Write your answer to each question in the space provided.
- Do **not** use an erasable pen or correction fluid.
- Do **not** write on any bar codes.
- You should use a calculator where appropriate.
- You must show all necessary working clearly; no marks will be given for unsupported answers from a calculator.
- Give non-exact numerical answers correct to 3 significant figures, or 1 decimal place for angles in degrees, unless a different level of accuracy is specified in the question.

INFORMATION

- The total mark for this paper is 80.
- The number of marks for each question or part question is shown in brackets [].

This document has **16** pages. Blank pages are indicated.

Mathematical Formulae**1. ALGEBRA***Quadratic Equation*For the equation $ax^2 + bx + c = 0$

$$x = \frac{-b \pm \sqrt{b^2 - 4ac}}{2a}$$

Binomial Theorem

$$(a+b)^n = a^n + \binom{n}{1}a^{n-1}b + \binom{n}{2}a^{n-2}b^2 + \dots + \binom{n}{r}a^{n-r}b^r + \dots + b^n,$$

where n is a positive integer and $\binom{n}{r} = \frac{n!}{(n-r)!r!}$

Arithmetic series

$$\begin{aligned} u_n &= a + (n-1)d \\ S_n &= \frac{1}{2}n(a+l) = \frac{1}{2}n\{2a + (n-1)d\} \end{aligned}$$

Geometric series

$$\begin{aligned} u_n &= ar^{n-1} \\ S_n &= \frac{a(1-r^n)}{1-r} \quad (r \neq 1) \\ S_\infty &= \frac{a}{1-r} \quad (|r| < 1) \end{aligned}$$

2. TRIGONOMETRY*Identities*

$$\begin{aligned} \sin^2 A + \cos^2 A &= 1 \\ \sec^2 A &= 1 + \tan^2 A \\ \operatorname{cosec}^2 A &= 1 + \cot^2 A \end{aligned}$$

Formulae for ΔABC

$$\begin{aligned} \frac{a}{\sin A} &= \frac{b}{\sin B} = \frac{c}{\sin C} \\ a^2 &= b^2 + c^2 - 2bc \cos A \\ \Delta &= \frac{1}{2}bc \sin A \end{aligned}$$


3**1** Sb e

$$xy = 3$$

$$x^4y^5 = 6$$

[3]

- 2** (a) On th a~~s~~ s b low , sk tch th g aph $y = \frac{1}{5}(x - 2)(x - 4)(x + 5)$, sh ig th co d n tes th p n s wh re th g aph eets th co d n te a~~s~~.



[2]

- (b) Expl ain wh~~y~~ sk tch ip rt (a) carib s ed os b v $(x - 2)(x - 4)(x + 5) \leq 0$ [1]

- (c) Hen e sb e $(x - 2)(x - 4)(x + 5) \leq 0$ [1]

3 Functions and their graphs

$$\begin{aligned}g(x) &= 2 + 4 \ln x \quad \text{for } x > 0 \\h(x) &= x^2 + 4 \quad \text{for } x > 0\end{aligned}$$

(a) Find g^{-1} , stating its domain and range.

[4]


(b) Solve $g(x) = 0$

[3]

- (c) Sb &
- $g'(x) = h'(x)$
- .

[3]

- 4 On the axes below, sketch the graph of $y = 2 \sin \frac{3}{2}x - 1$ for $0 \leq x \leq 60^\circ$, showing the coordinates of the points where the graph intersects the coordinate axes. [4]



- 5 (a) A 6 character password should be self contained following characters.

letters	A	B	E	F
m b rs	5	8	9	
sym b s	*		\$	

Each character may be selected once in a password.

Find how many different 6 character password that may be chosen if

- (i) there are no restrictions,

[1]

- (ii) the password does not start with m b s in that order,

[2]

- (iii) the password does not start with sym b .

[2]

- (b) An examination consists of a section A, containing 10 short questions, and a section B containing 5 long questions. Candidates are required to answer 6 questions from section A and 3 questions from section B.

Find the number of different selections of 6 questions that can be made if

- (i) there are no other restrictions,

[2]

- (ii) candidates must answer the first 2 questions in section A and the first question in section B.

[2]

- 6 A particle P travels in a straight line such that, t s after passing through a fixed point O , its velocity $v \text{ ms}^{-1}$ is given by $v = \left(e^{\frac{t^2}{8}} - 4 \right)^3$.

(a) Find the speed of P at O . [1]

(b) Find the value of t for which P is instantaneously at rest. [2]

(c) Find the acceleration of P when $t = 1$. [4]

- 7 Variables x and y are such that when y is plotted against x^2 , a straight line graph passing through the points (3, 1) and (4, 2) is obtained.


(a) Give in what $y = Ab^{x^2}$, find the values of each of the constants A and b . [4]

(b) Find the values of y when $x = 5$ [2]

(c) Find the positive values of x when $y = 2$ [2]

10

8



The diagram shows a circle, centre O , radius r cm. The points A and B lie on the circle such that angle $AOB = 2\theta$ radians.


- (a) Given that the perimeter of the shaded sector is 10 cm, where $r = \frac{10}{\theta + \sin\theta}$. [3]

11

- (b) Give the radius θ canary, find the area of $\frac{dr}{d\theta}$ when $\theta = \frac{\pi}{6}$.

[4]

9



It is also given that $\overrightarrow{AB} = 4\mathbf{a}$, $\overrightarrow{BC} = \mathbf{b}$ and $\overrightarrow{DC} = 7\mathbf{a}$. The lines AC and DB intersect at the point X .

Find in terms of \mathbf{a} and \mathbf{b} ,

(a) \overrightarrow{DB} ,

[1]

(b) \overrightarrow{DA} .

[1]

Given that $\overrightarrow{AX} = \lambda \overrightarrow{AC}$ find in terms of \mathbf{a} , \mathbf{b} and λ ,

(c) \overrightarrow{AX} ,

[1]

(d) \overrightarrow{DX} .

[2]

13


Given that $\overrightarrow{DX} = \mu \overrightarrow{DB}$,

(e) find the value of λ and μ .

[4]

14

- 10 (a) (i) Sketch the graph of $y = e^x - 5$ on the axes below, showing the exact coordinates where the graph intersects the axes.



[3]

- (ii) Find the range of values of k for which the equation $e^x - 5 = k$ has no solutions. [1]

15

- (b) Simplify $\log_a \sqrt{2} + \log_a 8 + \log_a \left(\frac{1}{2}\right)$, giving your answer in the form $p \log_a 2$ where p is a constant. [2]

- (c) Solve the equation $\log_3 x - \log_9 4x = 1$. [4]

Question 11 is printed on the next page.

11 (a) (i) Show that $\frac{\operatorname{cosec} \theta}{\operatorname{cosec} \theta - \sin \theta} = \sec^2 \theta$. [3]

(ii) Hence solve $\frac{2 \operatorname{cosec} \phi}{\operatorname{cosec} \phi - \sin \phi} = 8$ for $0 < \phi < \theta$. [3]

(b) Solve $\sqrt{3} \tan\left(x + \frac{\pi}{4}\right) = 1$ for $0 < x < 2\pi$, giving answers in terms of π . [3]